

You are scheduled to receive a SEDATED FLEXIBLE SIGMOIDOSCOPY EXAM.

Please read all of the following information carefully and ahead of time so that you can be adequately prepared for this procedure.

Do not hesitate to contact your doctor or call (858) 939-6531 with any questions or concerns.

Your Appointment Information	Your	Appointme	ent Infor	mation:
------------------------------	------	-----------	-----------	---------

Date:	
Time:	
Physician:	

Sharp Rees-Stealy San Diego Department of Gastroenterology

# WHAT IS A SEDATED FLEXIBLE SIGMOIDOSCOPY?

A sedated flexible sigmoidoscopy is a visual examination used to evaluate the colon. During a flexible sigmoidoscopy exam, a thin, flexible tube is inserted into the rectum. A small camera at the tip of the tube allows your doctor to view the inside of the rectum and most of the sigmoid colon. An intravenous line – or "IV" – will be placed in your arm to deliver fluids and sedation medications during the procedure for your comfort.

We want your sedated flexible sigmoidoscopy exam to be as comfortable and successful as possible and your attention and adherence to these guidelines is vital to this success. Please read this entire information packet and follow all of the instructions without skipping or altering any of the steps.

HOW TO PREPARE FOR YOUR FLEXIBLE SIGMOIDOSCOPY

	Purchase preparation materials at the Sharp Rees-Stealy Pharmacy prior to leaving the medical center.  • You will need to purchase one bottle of Magnesium Citrate and two Fleet Ready-to-Use Enemas to cleanse the colon in preparation for your sigmoidoscopy exam.				
	Purchase clear liquids – you cannot eat solid food after 2 p.m. the day prior to your procedure.				
	<ul> <li>Acceptable clear liquids include the following (<u>avoid red- or purple colored liquids</u>):</li> </ul>				
	<ul> <li>Strained fruit juices without pulp (such as apple, white grape and lemonade) – no purple or red varieties</li> <li>Water</li> <li>Clear broth or bouillon</li> <li>Coffee or tea (without milk, cream or non-dairy creamer)</li> <li>Gatorade/sports drinks – no purple or red varieties</li> <li>Carbonated and non-carbonated soft drinks – no purple or red varieties</li> <li>Plain Jell-O/gelatin dessert (no added fruit or toppings) – no purple or red varieties</li> <li>Popsicles – no purple or red varieties</li> </ul>				
	Secure a responsible companion to accompany you to your sigmoidoscopy.				

- As mentioned above, to ensure your comfort, IV sedation will be given to you during your flexible sigmoidoscopy. You must secure a responsible companion, family member or friend,18 years of age or older, to escort you to your procedure, remain on site during the procedure, and to drive you home or to accompany you home via transit services.
- If you do not have an escort with you upon arrival for your flexible sigmoidoscopy, your procedure will be cancelled. It is very important that you understand that because of the medications given during the exam, you will not be able to drive until the day following your procedure.

## FLEXIBLE SIGMOIDOSCOPY PREPARATION INSTRUCTIONS

Excellent preparation is crucial to assure proper cleansing of your colon, allowing for an accurate and effective sigmoidoscopy. Please follow these instructions fully and do not hesitate to contact your doctor's office if you have any questions or concerns.

You can create a personalized preparation schedule below by noting the dates and times of each preparation step as related to your sigmoidoscopy appointment time.

These preparations begin the day before your procedure. Remember, failure to follow these directions may result in the rescheduling of your procedure to a later date.

Please note that your body loses significant amounts of fluid during bowel preparation. To prevent dehydration, it is important to supplement that fluid loss with clear liquids (see the list of acceptable clear liquids earlier in this packet). Make a conscious effort to drink clear liquids as much as you can. However, please remember that <u>you may not</u> eat or drink anything for four hours before your procedure.


<b>(</b>	1 DAY PRIO	OR TO YOUR FLEXIBLE SIGMOIDOSCOPY
	•	Eat breakfast, a generous lunch and anything else you want before 2 p.m.
<b>(</b>		OR TO YOUR FLEXIBLE SIGMOIDOSCOPY, AT 2 P.M/TIME
	•	Stop eating all solid food. Do not eat anything until after your procedure tomorrow. You may drink as many clear liquids as you want until four hours prior to your exam (see the list of acceptable clear liquids earlier in this packet).
<b>(</b>		OR TO YOUR FLEXIBLE SIGMOIDOSCOPY, BETWEEN 6 – 7 P.M/TIME
	•	Drink the entire bottle of Magnesium Citrate and continue to drink clear liquids.
<b>(</b>		PRIOR TO YOUR FLEXIBLE SIGMOIDOSCOPY/TIME

 Do not take anything by mouth, including liquids – clear or other – for four hours prior to your procedure. Again, <u>do not eat or drink anything</u> in the four hours before your scheduled procedure time.

(1)	2 HOURS	PRIOR TO Y	OUR FLEXIBLE SIGMOIDOSCOPY
	DATE	/TIME	

- Use the two Fleet Ready-to-Use Enemas.
  - Insert the first enema rectally, hold it for five minutes and expel.
  - Insert the second enema rectally, hold it for five minutes and expel.

# DIRECTIONS TO SHARP REES-STEALY SAN DIEGO 2929 Health Center Drive, San Diego, CA 92123



### Directions to Sharp Rees-Stealy San Diego Medical Center from I-163 traveling south

- Exit Genesee Avenue, turn left/east
- Turn left on Health Center Drive
- Pass Vista Hill Avenue
- The medical center is on the right, just before you get to the parking structure
- Turn right at the entrance to Sharp Rees-Stealy San Diego Medical Center
- Turn right at your first opportunity
- Patient parking is located in the surface lot south of the SRS building

#### Directions to Sharp Rees-Stealy Medical Center from I-163 traveling north

- Exit Genesee Avenue, turn right/east
- Turn left on Health Center Drive
- Pass Vista Hill Avenue
- The medical center is on the right, just before you get to the parking structure
- Turn right at the entrance to Sharp Rees-Stealy San Diego Medical Center
- Turn right at your first opportunity
- Patient parking is located in the surface lot south of the SRS building

#### Directions to Sharp Rees-Stealy San Diego Medical Center from I-805 traveling north

- Exit Mesa College Drive/Kearny Villa Road, turn left/west
- Turn left on Health Center Drive
- Pass Frost Street
- The medical center is on the left, just past the parking structure
- Turn left at the entrance to Sharp Rees-Stealy San Diego Medical Center
- Turn right at your first opportunity
- Patient parking is located in the surface lot south of the SRS building

NOTE: If traveling south from I-805, there is no south-bound exit from the I-805 freeway. It is best to transfer to I-163, and then follow those directions above to the medical