

ECCO guidelines on therapeutics in ulcerative colitis: medical treatment.

Guideline ID: 3289

Published: 2022 Jan 28

European Crohn's and Colitis Organisation (ECCO)

Raine T, Bonovas S, Burisch J, et al. ECCO guidelines on therapeutics in ulcerative colitis: medical treatment. J Crohns Colitis. 2022 Jan 28;16(1):2-17. [162 references] [PubMed](#)

 [View Original Guideline](#)

Overview

Guideline Objective

To provide recommendations on medical treatment for adults with ulcerative colitis (UC)

Patient Population

Adults with UC

Recommendations

Recommendation Statements

Major interventions covered in this guideline include:

- Guidance on use of pharmacological interventions (e.g., 5-aminosalicylates (5-ASA); topical, colonic-release, and systemic corticosteroids; immunomodulators;

anti-tumour necrosis factor (TNF) agents), based on disease severity and activity

Note: Full recommendation statements have not been provided because this guideline does not meet [EGT's systematic review of the evidence criteria](#). Refer to the [original guideline](#) for more information.

Evidence Rating Scheme

Refer to the original guideline documentation for more information.

Recommendation Rating Scheme

Refer to the original guideline documentation for more information.

Related Content

Supporting Documents

- [Supplementary Material](#).
 - [Supplementary File 1](#).
 - [Supplementary File 2](#).
 - [Supplementary File 3](#).

Implementation Tools

No implementation tools available.

Patient Education

No patient education materials available.

Disclaimer

If you desire to use content from the original clinical practice guideline cited herein, you must contact the guideline developer directly to obtain permission rights.

ECRI's Guideline Profiles are designed to provide information and assist decision-making. Variations in practice will inevitably, and appropriately, occur when clinicians take into account the needs and preferences of individual patients, available resources, and limitations unique to an institution or type of practice. Every healthcare professional using these Guideline Profiles is responsible for evaluating the appropriateness of applying them in a clinical setting.