European Stroke Organisation (ESO) guidelines on management of transient ischaemic attack.

Guideline ID: 2551 Published: 2021 Jun

European Stroke Organisation (ESO)

Fonseca AC, Merwick A, Dennis M, et al. European Stroke Organisation (ESO) guidelines on management of transient ischaemic attack. Eur Stroke J. 2021 Jun;6(2):V. [92 references] *PubMed*

View Original Guideline

Overview

Guideline Objective

To provide clinically useful evidence-based recommendations on approaches to triage, investigation, and secondary prevention, particularly in the acute phase following transient ischaemic attack (TIA)

Patient Population

Adults with a suspected or confirmed TIA

Recommendations

Recommendation Statements

Major interventions covered in this guideline include:

- Role of specialist, including timing and setting of patient review
- Guidance on use of risk prediction tools
- Diagnostic imaging, including magnetic resonance imaging (MRI), computed tomography (CT), CT perfusion, MR angiogram (MRA), and CT angiography (CTA)
- Use of antiplatelet monotherapy and dual antiplatelet therapy (DAPT)

Note: Full recommendation statements have not been provided because this guideline does not meet EGT's systematic review of the evidence criteria. Refer to the original guideline for more information.

Evidence Rating Scheme

Refer to the original guideline documentation for more information.

Recommendation Rating Scheme

Refer to the original guideline documentation for more information.

Related Content

Supporting Documents

- Supplemental Material.

Implementation Tools

No implementation tools available.

Patient Education

No patient education materials available.

Disclaimer

If you desire to use content from the original clinical practice guideline cited herein, you must contact the guideline developer directly to obtain permission rights.

ECRI's Guideline Profiles are designed to provide information and assist decision-making. Variations in practice will inevitably, and appropriately, occur when clinicians take into account the needs and preferences of individual patients, available resources, and limitations unique to an institution or type of practice. Every healthcare professional using these Guideline Profiles is responsible for evaluating the appropriateness of applying them in a clinical setting.

Copyright © 2025 ECRI