

Treatment of urogenital symptoms in individuals with a history of estrogen-dependent breast cancer: ACOG clinical consensus no. 2.

Guideline ID: 2795

Published: 2021 Dec

American College of Obstetricians and Gynecologists (ACOG)

Treatment of urogenital symptoms in individuals with a history of estrogen-dependent breast cancer: ACOG clinical consensus no. 2. Obstet Gynecol. 2021 Dec 1;138(6):950-960. [86 references] [PubMed](#)

[View Original Guideline](#)

Overview

Guideline Objective

To review the safety and efficacy of newer hormonal treatment options as well as nonhormonal modalities

Patient Population

Individuals with a history of estrogen-dependent breast cancer

Recommendations

Recommendation Statements

Major interventions covered in this guideline include:

- Nonhormonal methods for urogenital and vulvovaginal symptoms, including silicone-, polycarbophil-, and water-based lubricants; hyaluronic acid; polyacrylic acid; and vitamin E and D vaginal suppositories
- Hormonal treatment options, including low-dose vaginal estrogen, vaginal dehydroepiandrosterone (DHEA) or testosterone, and ospemifene/selective estrogen receptor modulators
- Shared decision making

Note: Full recommendation statements have not been provided because this guideline does not meet [EGT's systematic review of the evidence criteria](#). Refer to the [original guideline](#) for more information.

Evidence Rating Scheme

Refer to the original guideline documentation for more information.

Recommendation Rating Scheme

Refer to the original guideline documentation for more information.

Related Content

Supporting Documents

- [Appendix 1. Literature Search Strategy.](#)
- [Appendix 2. Evidence Map.](#)
- [Clinical Consensus Methodology](#); 2021 Sep.

Implementation Tools

No implementation tools available.

Patient Education

No patient education materials available.

Disclaimer

If you desire to use content from the original clinical practice guideline cited herein, you must contact the guideline developer directly to obtain permission rights.

ECRI's Guideline Profiles are designed to provide information and assist decision-making. Variations in practice will inevitably, and appropriately, occur when clinicians take into account the needs and preferences of individual patients, available resources, and limitations unique to an institution or type of practice. Every healthcare professional using these Guideline Profiles is responsible for evaluating the appropriateness of applying them in a clinical setting.