EASL clinical practice guidelines on non-invasive tests for evaluation of liver disease severity and prognosis – 2021 update.

Guideline ID: 2807 Published: 2021 Sep

European Association for the Study of the Liver (EASL)

European Association for the Study of the Liver. EASL clinical practice guidelines on non-invasive tests for evaluation of liver disease severity and prognosis – 2021 update. J Hepatol. 2021 Sep;75(3):659-689. [307 references] PubMed

View Original Guideline

Overview

Guideline Objective

To provide recommendations on the use of non-invasive tests for the evaluation of liver disease severity and prognosis

Patient Population

Adult patients with suspected or confirmed chronic liver disease

Recommendations

Recommendation Statements

Major interventions covered in this guideline include:

- Detection accuracy and use of non-invasive ventilation (NIV) scores, serum markers, liver stiffness, imaging methods, and liver biopsy for diagnostic and prognostic assessment

Note: Full recommendation statements have not been provided because this guideline does not meet EGT's systematic review of the evidence criteria. Refer to the original guideline for more information.

Evidence Rating Scheme

Refer to the original guideline documentation for more information.

Recommendation Rating Scheme

Refer to the original guideline documentation for more information.

Related Content

Supporting Documents

- Supplementary Data: Recommendations that Remained Unchanged as in the Previous EASL CPGs.
- Supplementary Data: Disclosure Form.
- Clinical Practice Guidelines of the European Association for the Study of the Liver – Advancing Methodology but Preserving Practicability; 2019 Jan 1.

Implementation Tools

No implementation tools available.

Patient Education

No patient education materials available.

Disclaimer

If you desire to use content from the original clinical practice guideline cited herein, you must contact the guideline developer directly to obtain permission rights.

ECRI's Guideline Profiles are designed to provide information and assist decision-making. Variations in practice will inevitably, and appropriately, occur when clinicians take into account the needs and preferences of individual patients, available resources, and limitations unique to an institution or type of practice. Every healthcare professional using these Guideline Profiles is responsible for evaluating the appropriateness of applying them in a clinical setting.

Copyright © 2025 ECRI