

PA007ITRI06
VL7516/ HP8799

For more information, visit us at www.lillyindia.co.in

Insulin Lispro Injection I.P. (r-DNA origin) Humalog®

(100 IU/mL, 3 mL solution for injection in cartridge/
solution for injection in a pre-filled pen)

[Monocomponent Insulin lispro, recombinant DNA origin]

1. NAME OF THE MEDICINAL PRODUCT

Humalog® 100 IU/mL, solution for injection in cartridge
Humalog KwikPen® 100 IU/mL, solution for injection in a pre-filled pen

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each mL of Insulin Lispro contains: 100 IU (equivalent to 3.5 mg) Insulin Lispro I.P. (r-DNA origin) as active ingredient, 16 mg Glycerol I.P. as tonicity modifier, 3.15 mg Metacresol USP as preservative/stabilizer, Zinc Oxide I.P. q.s. as stabilizer, 1.88 mg Dibasic Sodium Phosphate I.P. as buffering agent, Hydrochloric Acid I.P. (10 %) q.s. and Sodium Hydroxide I.P. (10 %) q.s. for pH adjustment, Water for Injection I.P. q.s. 1mL.

Cartridge

Each cartridge contains 300 units of insulin lispro in 3 ml solution.

KwikPen

Each pre-filled pen contains 300 units of insulin lispro in 3 ml solution. Each KwikPen delivers 1-60 units in steps of 1 unit.

3. PHARMACEUTICAL FORM

Solution for injection. Humalog® is a sterile, clear, colourless, aqueous solution.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Humalog® is indicated for the treatment of patients with diabetes mellitus who require insulin for the maintenance of normal glucose homeostasis. Humalog® is also indicated for the initial stabilisation of diabetes mellitus.

4.2 Posology and method of administration

Posology

The dosage should be determined by the physician, according to the requirement of the patient.

Humalog® may be given shortly before meals. When necessary Humalog® can be given soon after meals.

Humalog® takes effect rapidly and has a shorter duration of activity (2 to 5 hours) given subcutaneously as compared with soluble insulin. This rapid onset of activity allows a Humalog® injection (or, in the case of administration by continuous subcutaneous infusion, a Humalog® bolus) to be given very close to mealtime.

The time course of action of any insulin may vary considerably in different individuals or at different times in the same individual. The faster onset of action compared to soluble human insulin is maintained regardless of injection site. As with all insulin preparations, the duration of action of Humalog® is dependent on dose, site of injection, blood supply, temperature, and physical activity.

Humalog® can be used in conjunction with a longer-acting insulin or oral sulphonylurea agents, on the advice of a physician.

Humalog KwikPens

Humalog KwikPen® is available in two strengths. The Humalog 100 units/ml KwikPen (and Humalog 200 units/ml KwikPen, see separate SmPC) delivers 1 – 60 units in steps of 1 unit in a single injection. **The number of insulin units is shown in the dose window of the pen regardless of strength and no dose conversion should be done when transferring a patient to a new strength or to a pen with a different dose step.**

Special populations

Renal impairment

Insulin requirements may be reduced in the presence of renal impairment.

Hepatic impairment

Insulin requirements may be reduced in patients with hepatic impairment due to reduced capacity for gluconeogenesis and reduced insulin breakdown; however, in patients with chronic hepatic impairment, an increase in insulin resistance may lead to increased insulin requirements.

Paediatric population

Humalog® can be used in adolescents and children (see section 5.1)

Method of administration

Humalog® preparations should be given by subcutaneous injection.

The KwikPen is only suitable for subcutaneous injections. Humalog® in cartridges is only suitable for subcutaneous injections from a Lilly reusable pen or compatible pump systems for continuous subcutaneous insulin infusion (CSII).

Subcutaneous administration should be in the upper arms, thighs, buttocks, or abdomen. Use of injection sites should be rotated so that the same site is not used more than approximately once a month.

When administered subcutaneously care should be taken when injecting Humalog® to ensure that a blood vessel has not been entered. After injection, the site of injection should not be massaged. Patients must be educated to use the proper injection techniques.

Use of Humalog® in an insulin infusion pump

For subcutaneous injection of Humalog® using a continuous infusion pump, you may fill the pump reservoir from a Humalog® 100 units/ml vial. Some pumps are compatible with cartridges that can be inserted intact into the pump.

Only certain CE-marked insulin infusion pumps may be used to infuse insulin lispro. Before infusing insulin lispro, the pump manufacturer's instructions should be studied to ascertain the suitability for the particular pump. Use the correct reservoir and catheter for the pump. When filling the pump reservoir avoid damaging it by using the correct needle length on the filling system. The infusion set (tubing and cannula) should be changed in accordance with the instructions in the product information supplied with the infusion set. In the event of a hypoglycaemic episode, the infusion should be stopped until the episode is resolved. If repeated or severe low blood glucose levels occur consider the need to reduce or stop an insulin infusion. A pump malfunction or obstruction of the infusion set can result in a rapid rise in glucose levels.

If an interruption to insulin flow is suspected, follow the instructions in the pump product literature. When used with an insulin infusion pump, Humalog should not be mixed with any other insulin.

Intravenous administration of insulin

If necessary, Humalog® may also be administered intravenously, for example: for the control of blood glucose levels during ketoacidosis, acute illnesses or during intra and post operative periods.

Intravenous injection of insulin lispro should be carried out following normal clinical practise for intravenous injections, for example by an intravenous bolus or by an infusion system. Frequent monitoring of the blood glucose levels is required.

Infusion systems at concentrations from 0.1 units/ml to 1.0 units/ml insulin lispro in 0.9% sodium chloride or 5% dextrose are stable at room temperature for 48 hours. It is recommended that the system is primed before starting the infusion to the patient.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Hypoglycaemia.

4.4 Special warnings and precautions for use

Traceability

In order to improve the traceability of biological medicinal products, the name and the batch number of the administered product should be clearly recorded.

Transferring a patient to another type or brand of insulin

Transferring a patient to another type or brand of insulin should be done under strict medical supervision. Changes in strength, brand (manufacturer), type (regular/soluble, NPH/isophane etc.), species (animal, human, human insulin analogue), and/or method of manufacture (recombinant DNA versus animal-source insulin) may result in the need for a change in dosage. For fast-acting insulins, any patient also on basal insulin must optimise dosage of both insulins to obtain glucose control across the whole day, particularly nocturnal/fasting glucose control.

Hypoglycaemia and hyperglycaemia

Conditions which may make the early warning symptoms of hypoglycaemia different or less pronounced include long duration of diabetes, intensified insulin therapy, diabetic nerve disease or medications such as beta-blockers.

A few patients who have experienced hypoglycaemic reactions after transfer from animal-source insulin to human insulin have reported that the early warning symptoms of hypoglycaemia were less pronounced or different from those experienced with their previous insulin. Uncorrected hypoglycaemic or hyperglycaemic reactions can cause loss of consciousness, coma, or death. The use of dosages which are inadequate or discontinuation of treatment, especially in insulin dependent diabetics, may lead to hyperglycaemia and diabetic ketoacidosis; conditions which are potentially lethal.

Insulin requirements and dosage adjustment

Insulin requirements may be increased during illness or emotional disturbances.

Adjustment of dosage may also be necessary if patients undertake increased physical activity or change their usual diet. Exercise taken immediately after a meal may increase the risk of hypoglycaemia. A consequence of the pharmacodynamics of rapid-acting insulin analogues is that if hypoglycaemia occurs, it may occur earlier after an injection when compared with soluble human insulin.

Combination of Humalog with pioglitazone:

Cases of cardiac failure have been reported when pioglitazone was used in combination with insulin, especially in patients with risk factors for development of cardiac heart failure. This should be kept in mind, if treatment with the combination of pioglitazone and Humalog® is considered. If the combination is used, patients should be observed for signs and symptoms of heart failure, weight gain and oedema. Pioglitazone should be discontinued, if any deterioration in cardiac symptoms occurs.

Avoidance of medication errors

Patients must be instructed to always check the insulin label before each injection to avoid accidental mix-ups between the two different strengths of Humalog KwikPen® as well as other insulin products.

Patients must visually verify the dialled units on the dose counter of the pen. Therefore, the requirement for patients to self-inject is that they can read the dose counter on the pen. Patients who are blind or have poor vision must be instructed to always get help/assistance from another person who has good vision and is trained in using the insulin device.

Excipients

This medicinal product contains less than 1 mmol sodium (23 mg) per dose, i.e., essentially "sodium-free".

4.5 Interaction with other medicinal products and other forms of interaction

Insulin requirements may be increased by medicinal products with hyperglycaemic activity, such as oral contraceptives, corticosteroids, or thyroid replacement therapy, danazol, beta2 stimulants (such as ritodrine, salbutamol, terbutaline).

Insulin requirements may be reduced in the presence of medicinal products with hypoglycaemic activity, such as oral hypoglycaemics, salicylates (for example, acetylsalicylic acid), sulpha antibiotics, certain antidepressants (monoamine oxidase inhibitors, selective serotonin reuptake inhibitors), certain angiotensin converting enzyme inhibitors (captopril, enalapril), angiotensin II receptor blockers, beta-blockers, octreotide or alcohol.

The physician should be consulted when using other medications in addition to Humalog® (see section 4.4).

4.6 Fertility, pregnancy and lactation

Pregnancy

Data on a large number of exposed pregnancies do not indicate any adverse effect of insulin lispro on pregnancy or on the health of the foetus/newborn.

It is essential to maintain good control of the insulin-treated (insulin-dependent or gestational diabetes) patient throughout pregnancy. Insulin requirements usually fall during the first trimester and increase during the second and third trimesters. Patients with diabetes should be advised to inform their doctor if they are pregnant or are contemplating pregnancy. Careful monitoring of glucose control, as well as general health, is essential in pregnant patients with diabetes.

Breast-feeding

Patients with diabetes who are breast-feeding may require adjustments in insulin dose, diet or both.

Fertility

Insulin lispro did not induce fertility impairment in animal studies (see section 5.3).

4.7 Effects on ability to drive and use machines

The patient's ability to concentrate and react may be impaired as a result of hypoglycaemia. This may constitute a risk in situations where these abilities are of special importance (e.g. driving a car or operating machinery).

Patients should be advised to take precautions to avoid hypoglycaemia whilst driving, this is particularly important in those who have reduced or absent awareness of the warning signs of hypoglycaemia or have frequent episodes of hypoglycaemia. The advisability of driving should be considered in these circumstances.

4.8 Undesirable effects

Summary of safety profile

Hypoglycaemia is the most frequent undesirable effect of insulin therapy that a patient with diabetes may suffer. Severe hypoglycaemia may lead to loss of consciousness, and in extreme cases, death. No specific frequency for hypoglycaemia is presented, since hypoglycaemia is a result of both the insulin dose and other factors e.g. a patient's level of diet and exercise.

Tabulated list of adverse reactions

The following related adverse reactions from clinical trials are listed below as MedDRA preferred term by system organ class and in order of decreasing incidence (very common: ≥1/10; common: ≥1/100 to <1/10; uncommon: ≥1/1,000 to <1/100; rare: ≥1/10,000 to <1/1,000; very rare: <1/10,000).

Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

MedDRA system organ classes	Very common	Common	Uncommon	Rare	Very rare
Immune system disorders					
Local allergy		X			
Systemic allergy				X	
Skin and subcutaneous tissue disorders					
Lipodystrophy			X		

PPD Information Box		ALRP Information Box
Technical Information:	<input checked="" type="checkbox"/> BLACK <input type="checkbox"/> DIECUT	Translations of Variable Data Lot: N/A Exp Date: N/A Mfg Date: N/A Price: N/A GTIN: N/A Serial Number : N/A
Layout Name	Previous Item Code (to be destroyed)	Variable Barcode Information
SESPA200A00	PA007ITRI05	N/A

Description of selected adverse reactions

Local allergy

Local allergy in patients is common. Redness, swelling, and itching can occur at the site of insulin injection. This condition usually resolves in a few days to a few weeks. In some instances, this condition may be related to factors other than insulin, such as irritants in the skin cleansing agent or poor injection technique.

Systemic allergy

Systemic allergy, which is rare but potentially more serious, is a generalized allergy to insulin. It may cause a rash over the whole body, shortness of breath, wheezing, reduction in blood pressure, fast pulse, or sweating. Severe cases of generalised allergy may be life-threatening.

Lipodystrophy

Lipodystrophy at the injection site is uncommon.

Oedema

Cases of oedema have been reported with insulin therapy, particularly if previous poor metabolic control is improved by intensified insulin therapy.

4.9 Overdose

Insulins have no specific overdose definitions because serum glucose concentrations are a result of complex interactions between insulin levels, glucose availability and other metabolic processes.

Hypoglycaemia may occur as a result of an excess of insulin activity relative to food intake and energy expenditure.

Hypoglycaemia may be associated with listlessness, confusion, palpitations, headache, sweating and vomiting. Mild hypoglycaemic episodes will respond to oral administration of glucose or other sugar or saccharated products. Correction of moderately severe hypoglycaemia can be accomplished by intramuscular or subcutaneous administration of glucagon, followed by oral carbohydrate when the patient recovers sufficiently. Patients who fail to respond to glucagon must be given glucose solution intravenously. If the patient is comatose, glucagon should be administered intramuscularly or subcutaneously. However, glucose solution must be given intravenously if glucagon is not available or if the patient fails to respond to glucagon. The patient should be given a meal as soon as consciousness is recovered. Sustained carbohydrate intake and observation may be necessary because hypoglycaemia may recur after apparent clinical recovery.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group : Drugs used in diabetes, insulins and analogues for injection, fast-acting

ATC code: A10A B04


The primary activity of insulin lispro is the regulation of glucose metabolism.

In addition, insulins have several anabolic and anti-catabolic actions on a variety of different tissues. Within muscle tissue this includes increasing glycogen, fatty acid, glycerol and protein synthesis and amino acid uptake, while decreasing glycogenolysis, gluconeogenesis, ketogenesis, lipolysis, protein catabolism and amino acid output.

Insulin lispro has a rapid onset of action (approximately 15 minutes), thus allowing it to be given closer to a meal (within zero to 15 minutes of the meal) when compared to soluble insulin (30 to 45 minutes before). Insulin lispro takes effect rapidly and has a shorter duration of activity (2 to 5 hours) when compared to soluble insulin.

Clinical trials in patients with type 1 and type 2 diabetes have demonstrated reduced postprandial hyperglycaemia with insulin lispro compared to soluble human insulin.

As with all insulin preparations, the time course of insulin lispro action may vary in different individuals or at different times in the same individual and is dependent on dose, site of injection, blood supply, temperature and physical activity. The typical activity profile following subcutaneous injection is illustrated below.



The above representation reflects the relative amount of glucose over time required to maintain the subject's whole blood glucose concentrations near fasting levels and is an indicator of the effect of these insulins on glucose metabolism over time.

Clinical trials have been performed in children (61 patients aged 2 to 11) and children and adolescents (481 patients aged 9 to 19 years), comparing insulin lispro to human soluble insulin. The pharmacodynamic profile of insulin lispro in children is similar to that seen in adults.

When used in subcutaneous infusion pumps, treatment with insulin lispro has been shown to result in lower glycosylated haemoglobin levels compared to soluble insulin. In a double-blind, crossover study, the reduction in glycosylated haemoglobin levels after 12 weeks dosing was 0.37 percentage points with insulin lispro, compared to 0.03 percentage points for soluble insulin ($p = 0.004$).

In patients with type 2 diabetes on maximum doses of sulphonyl urea agents, studies have shown that the addition of insulin lispro significantly reduces HbA1c compared to sulphonyl urea alone. The reduction of HbA1c would also be expected with other insulin products e.g. soluble or isophane insulins.

Clinical trials in patients with type 1 and type 2 diabetes have demonstrated a reduced number of episodes of nocturnal hypoglycaemia with insulin lispro compared to soluble human insulin. In some studies, reduction of nocturnal hypoglycaemia was associated with increased episodes of daytime hypoglycaemia.

The glucodynamic response to insulin lispro is not affected by renal or hepatic function impairment. Glucodynamic differences between insulin lispro and soluble human insulin, as measured during a glucose clamp procedure, were maintained over a wide range of renal function.

Insulin lispro has been shown to be equipotent to human insulin on a molar basis but its effect is more rapid and of a shorter duration.

5.2 Pharmacokinetic properties

The pharmacokinetics of insulin lispro reflect a compound that is rapidly absorbed, and achieves peak blood levels 30 to 70 minutes following subcutaneous injection. When considering the clinical relevance of these kinetics, it is more appropriate to examine the glucose utilisation curves (as discussed in 5.1).

Insulin lispro maintains more rapid absorption when compared to soluble human insulin in patients with renal impairment. In patients with type 2 diabetes over a wide range of renal function the pharmacokinetic differences between insulin lispro and soluble human insulin were generally maintained and shown to be independent of renal function. Insulin lispro maintains more rapid absorption and elimination when compared to soluble human insulin in patients with hepatic impairment.

5.3 Preclinical safety data

In vitro tests, including binding to insulin receptor sites and effects on growing cells, insulin lispro behaved in a manner that closely resembled human insulin. Studies also demonstrate that the dissociation of binding to the insulin receptor of insulin lispro is equivalent to human insulin. Acute, one month and twelve month toxicology studies produced no significant toxicity findings.

Insulin lispro did not induce fertility impairment, embryotoxicity or teratogenicity in animal studies.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Glycerol

m-Cresol

Zinc Oxide

Dibasic Sodium Phosphate

Water for Injection

Hydrochloric acid and sodium hydroxide maybe used to adjust pH

6.2 Incompatibilities

Cartridge and KwikPen

These medicinal products should not be mixed with any other insulin or any other medicinal product.

6.3 Shelf life

Before use: 3 years

After first use/ after cartridge insertion: 28 days

6.4 Special precautions for storage

Do not freeze. Do not expose to excessive heat or direct sunlight.

Before use

Store at 2°C - 8°C (in a refrigerator).

After first use / after cartridge insertion

Cartridge

Store below 30°C. Do not refrigerate. The pen with the inserted cartridge should not be stored with the needle attached.

KwikPen

Store below 30°C. Do not refrigerate. The pre-filled pen should not be stored with the needle attached.

6.5 Nature and contents of container

Cartridge

The solution is contained in type I flint glass cartridges, sealed with butyl or halobutyl disc seals and plunger heads, and are secured with aluminium seals. Dimeticone or silicone emulsion may be used to treat the cartridge plungers, and/or the glass cartridges.

3 ml Cartridge: Packs of 5 or 10. Not all packs may be marketed

KwikPen

The solution is contained in type I flint glass cartridges, sealed with butyl or halobutyl disc seals and plunger heads and are secured with aluminium seals. Dimeticone or silicone emulsion may be used to treat the cartridge plunger, and/or the glass cartridge. The 3 ml cartridges are sealed in a disposable pen injector, called the "KwikPen". Needles are not included.

3 ml KwikPen: Packs of 5 or a multipack of 10 (2 packs of 5). Not all packs may be marketed

6.6 Special precautions for disposal and other handling

Instructions for use and handling

To prevent the possible transmission of disease, each cartridge or pen must be used by one patient only, even if the needle on the delivery device is changed.

The Humalog® solution should be clear and colourless. Humalog® should not be used if it appears cloudy, thickened, or slightly coloured or if solid particles are visible.

Preparing a dose

Cartridge

Humalog® cartridges are to be used with a Lilly reusable insulin pen and should not be used with any other reusable pen as the dosing accuracy has not been established with other pens. The instructions with each individual pen must be followed for loading the cartridge, attaching the needle and administering the insulin injection.

KwikPen

Before using the pre-filled pen the user manual included in the package leaflet must be read carefully.

The pre-filled pen has to be used as recommended in the user manual.

Pens should not be used if any part looks broken or damaged.

Injecting a dose

If using a pre-filled or reusable pen refer to the detailed instructions for preparing the pen and injecting the dose, the following is a general description.

1. Wash your hands.
2. Choose a site for injection.
3. Clean the skin as instructed.
4. Stabilise the skin by spreading it or pinching up a large area. Insert the needle and inject as instructed.
5. Pull the needle out and apply gentle pressure over the injection site for several seconds. Do not rub the area.
6. Dispose of the syringe and needle safely. For an injection device use the outer needle cap, unscrew the needle and dispose of it safely
7. Use of the injection sites should be rotated so that the same is not used more than approximately once a month.

Any unused product or waste material should be disposed of in accordance with local requirements.

Manufactured by:

M/s. Eli Lilly Italia S.P.A., via Gramsci 731-733, 50019 Sesto Fiorentino (FI), Italy

*Marketed By:

Eli Lilly and Company (India) Pvt. Ltd.

Plot No. 92, Sector-32, Gurgaon-122001, Haryana, India

Imported By:

Eli Lilly and Company (India) Pvt. Ltd., Bldg. No. 14, Gala No. 1 to 4, 1st Fl, Arihant Comm. Complex, Purna Bhiwandi, Maharashtra-421302

If you have any questions or complaints with your Humalog® cartridge, contact Lilly at Toll Free number 18001230021 or your healthcare professional for assistance

*Under license from registered trademark owners Eli Lilly and Company, USA

Humalog® cartridges are designed to be used only with Lilly Insulin delivery devices

PA007ITRI06

PPD Information Box		ALRP Information Box
Technical Information: ■ BLACK □ DIECUT		Translations of Variable Data Lot: N/A Exp Date: N/A Mfg Date: N/A Price: N/A GTIN: N/A Serial Number: N/A
Layout Name SESPA200A00	Previous Item Code (to be destroyed) PA007ITRI05	Variable Barcode Information N/A