

Home Care After Your Celiac Plexus Block or Neurolysis (Nerve Destruction)

What to Expect

You may notice a mild soreness in your throat. You may wish to gargle with lightly salted water or suck on hard candy.

You may feel full or have gas pains.

At Home

You **must** have an adult to take you home. You should not drive today. Do not make important decisions today.

Do not drink alcohol. Do not take tranquilizers or sleeping pills tonight.

Rest today. You may resume normal daily tasks tomorrow.

You may resume your routine medicines unless told otherwise. Your doctor will tell you when to restart your blood thinners.

Your doctor will let you know if you need to be on a clear liquid diet, otherwise, start with a light diet the day of the procedure.

You may have some soreness and redness at the IV site for a day or two. You can relieve this by placing a warm, moist washcloth over the area. Please contact your primary doctor if this does not improve in a couple of days.

When to Call

- Fever above 100.4°F or 38°C for 2 readings taken 4 hours apart
- Pain that is new or more severe than the pain you had before the procedure
- New symptoms in your legs such as numbness, tingling or weakness
- Nausea, vomiting, abdominal fullness, or bloating
- Increasing stomach or back pain
- Fainting spells
- Blood-stained urine or stools
- Severe diarrhea

Who to Call

If you have questions or concerns within 24 hours of your procedure, please call the Procedure Center at the number below.

Doctor _____

Gastroenterology Procedures
Monday - Friday 7:00 a.m. - 5:00 p.m.
(608) 263-8094.

After clinic hours, call the paging operator at **(608) 262-0486** or **1-800-323-8942**. Ask for the Gastroenterology (GI) doctor on call. Leave your name and phone number with the area code. The doctor will call you back.