

Rectal Exam Under Anesthesia (With Pouch)

This is an exam of the rectum and ileal pouch while you are asleep under anesthesia.

Bowel Prep

You will need to do a bowel prep before the exam. The bowel prep is the MiraLAX prep.

MiraLAX Prep

You will need to purchase these items:

- MiraLAX 4.1 ounce/119 grams
- 2-32-ounce bottle of Gatorade or Powerade. (If diabetic, use Pedialyte)

The Day Before Surgery

You may eat a light lunch prior to 12:00 pm. No solid foods after 12:00 pm. You may have:

- Water, sparkling water or soda
- Broth
- Juice, without pulp (apple, grape)
- Popsicles
- Powerade
- Clear Jell-O® (no fruit, etc. in it) NO Jell-O® cups
- Coffee or tea, no creamer
- Gatorade®
- Crystal Light®

After 12:00 pm (noon), **do not** eat solid food. **Do not** drink juice with pulp, dairy products, or alcohol.

Day Before Surgery Steps

1. **2:00 pm:** In the early afternoon split the MiraLAX into the 2-32 bottles of chilled Gatorade or Powerade. (If diabetic, use Pedialyte)
2. **4:00 pm:** Drink the first 32-ounce bottle of the Gatorade/MiraLAX drink. Drink a glass full of the drink every 15 minutes until your stool is clear.

3. **6:00 pm:** If the stool is still not clear then drink the second 32-ounce Gatorade and MiraLAX mixture. Drink a glass full every 15 minutes. If stools clear, then you can stop drinking before you complete the second bottle. Continue to drink clear fluids until bedtime to prevent getting dehydrated.

4. **Before bed:** Shower.

The Day of Surgery

- Shower
- Take your medicines as you were told by the colorectal surgery team.
- Call if you have not gotten instructions.
- You may drink clear liquids until **4 hours** before you arrive for the procedure.

After the Exam

Your care after the exam will depend on the findings. If you had a dilation, you may be able to return to your normal routine in 1-2 days **or** you may need to follow more strict guidelines.

Activities

- Activity as tolerated.
- There may be restriction to activity depending what was done during the exam.
- Follow up will be determined by the surgeon.

Pain Relief

You may have pain after your exam. Your doctor may order pain pills for you.

When to Call the Doctor

- Problems having a bowel movement.
- Temperature by mouth over 100.4° F for two readings taken four hours apart.
- Pain not relieved by pain pills.
- Bleeding that does not stop after 10 minutes of applied pressure to the rectal area.

Phone Numbers

Digestive Health Center: **(608) 242-2800**

After hours, weekends or holidays the number will be answered by the paging operator. Ask for the doctor on call. Leave your name and phone number with area code. The doctor will call you back.

If you live out of the area, call **(855) 342-9900**.

If you are a patient receiving care at UnityPoint – Meriter, Swedish American or a health system outside of UW Health, please use the phone numbers provided in your discharge instructions for any questions or concerns.

Your health care team may have given you this information as part of your care. If so, please use it and call if you have any questions. If this information was not given to you as part of your care, please check with your doctor. This is not medical advice. This is not to be used for diagnosis or treatment of any medical condition. Because each person's health needs are different, you should talk with your doctor or others on your health care team when using this information. If you have an emergency, please call 911. Copyright © 4/2019 University of Wisconsin Hospitals and Clinics Authority. All rights reserved. Produced by the Department of Nursing. HF#4756.