

Going Home with a Peripheral Nerve Catheter

We will be giving you a local numbing medicine through a small catheter (tube) placed near the nerves that lead to the site of surgery. This medicine will help with your pain after surgery. You will likely have the catheter for 2-3 days. This handout will help to answer common questions.

The local pain medicine will not take away all your pain. You will likely still use some of the pain pills your doctor prescribed.

You will be getting a constant, set amount of numbing medicine every hour.

Most often, in about 10-15 hours, the intense numbness that you felt at first wears off and you may feel some pain.

When this occurs, take some of your pain medicine and press the bolus button on the front of your pump.

Most patients describe the treated area as feeling “fat” or numb.

For an upper extremity catheter, your fingers may feel “fat.” Still, you should be able to move your fingers and have some feeling in them. They just may not feel normal to you while the catheter is in place.

For a lower extremity catheter placed in the groin, your thigh should feel numb and weak. Nerve catheters placed in the groin should not cause you to lose feeling in your toes.

Sometimes we place nerve catheters in the back of the leg near your sciatic nerve. These catheters may cause foot and toe numbness.

The infusion should be off **for at least 2 hours** before you remove the catheter. You should have feeling back in the treated area.

Taking out the Catheter

1. Wash your hands with soap and water for 15-30 seconds.
2. Remove the dressing. Start with the tape holding the end of the catheter that connects to the pump tubing. Loosen the tape all the way down to where the catheter is inserted.
3. Gently pull the catheter out. It should come out smoothly. It may get hung up where there is glue. You can loosen up this glue so the catheter can move more easily if this is the case.
4. If it hurts to remove it or if you feel pain down your arm or leg while it is coming out, **stop and call the number listed** under “Who to Call.”
5. The tip at the end of the catheter should be silver, blue or black. If it is not, please call.
6. **Never cut the catheter!** Never disconnect the catheter from the tubing while it is in your leg, arm or shoulder.
7. Throw away the tubing, the medicine bag and catheter in the trash.

Guidelines

Do not drive while you have this medicine or while you take opioid pain medicine.

If we gave you medicine to help you relax or put you to sleep, do not make important decisions until the next day. You will need to have someone with you for the first 24 hours after you go home and for most of the time you get this medicine.

While bathing, do not get the catheter, pump, or wound wet. Sponge baths work best. You will need help bathing.

Cover the catheter site with a dressing. You do not need to change the dressing unless we tell you to. Fluid may leak around the catheter. This is normal and does not mean the medicine is not working.

Avoid contact with sharp surfaces or extreme heat or cold.

Arm and Upper Extremity Catheters

Keep your arm protected while you get this medicine. Keep it in a sling except while doing physical therapy.

If you had shoulder surgery, you may notice some shortness of breath, mainly when you lie down. This may be from the medicine. It will likely go away when the medicine is done. **Call your doctor and stop your pump if you are having shortness of breath.**

Leg and Lower Extremity Catheters

You are at risk for falling because you have a weak leg. If you have a nerve catheter near the groin, keep your knee in a locked knee brace, while walking, for the entire time you are getting this medicine.

You **must** use crutches or a walker while walking around. You will **not** be able to support your weight without these aids.

If you are a patient receiving care at UnityPoint – Meriter, Swedish American or a health system outside of UW Health, please use the phone numbers provided in your discharge instructions for any questions or concerns.

Your health care team may have given you this information as part of your care. If so, please use it and call if you have any questions. If this information was not given to you as part of your care, please check with your doctor. This is not medical advice. This is not to be used for diagnosis or treatment of any medical condition. Because each person's health needs are different, you should talk with your doctor or others on your health care team when using this information. If you have an emergency, please call 911. Copyright © 4/2020 University of Wisconsin Hospitals and Clinics Authority. All rights reserved. Produced by the Department of Nursing HF#6870

When to Call

- If you have questions or concerns about your catheter.
- If the catheter disconnects from the pump or falls out.
- If you have severe pain.
- If the skin around the catheter gets red, painful or appears infected.
- Symptoms of too much medicine (very rare):
 - Feel very sleepy
 - Have slurred speech
 - Tongue feels very thick
 - Have ringing in ears
 - Feel numbness or tingling around lips, tongue
 - Have metallic taste in mouth
 - Feel nervous or confused
- For serious health problems (seizures, heart symptoms, loss/decreased level of consciousness), please clamp the pump and **call 911** right away.

About the Pump

As you receive the medicine, the pump balloon will slowly become smaller.

MRI Tests

If you need an MRI, please tell your doctor you have a catheter and pump.

Who to Call

Ask for the resident on call for acute pain.
608-263-6400