

Getting Ready for an Enteroscopy (Push, Single or Double Balloon)

What is an Enteroscopy?

An enteroscopy (push endoscope) is an exam done to look further into the small intestine than a normal endoscope. During the exam a flexible tube with a light and camera is passed through your mouth, esophagus (the tube that connects the throat to the stomach), stomach, and into the small bowel. You may be sedated for the exam.

What is a Single Balloon Enteroscopy?

A single balloon enteroscopy is the same as above; but the flexible tube is fitted with another tube that has a balloon. The balloon can be inflated or deflated. This helps the doctor to pass further into the small intestine. You may be sedated for the exam.

What is a Double Balloon Enteroscopy?

A double balloon enteroscopy is the same as above; but the flexible tube is fitted with another tube that has 2 balloons that can be inflated or deflated. This helps the doctor to pass further into the small intestine. If the exam is being done to look at the lower part of the intestine, you will likely be sedated. If the exam is being done to look at the **upper** part of the intestine, it is always done with **general anesthesia**.

***Important – Please read this material 1 week in advance. Your procedure may be cancelled if you do not follow the instructions.**

Important to Know

1. Never stop taking blood thinning medicines without talking with your doctor first. They will give you instructions on how to safely stop these medicines. Call us if you have questions.

- a. **7 Days Before**

Stop taking the blood thinner Effient® (prasugrel).

- b. **5 Days Before**

Stop taking the blood thinners Coumadin® (warfarin) or Plavix® (clopidogrel).

- c. **2 Days Before**

Stop taking the blood thinners Pradaxa® (dabigatran), Xeralto® (rivaroxaban), Eliquis® (apixaban) and Brilinta® (ticagrelor).

2. Diabetes Medicines (oral and insulin) will need to be adjusted for the time you're not eating like you normally do. Please discuss this with your primary doctor.
3. **Plan ahead.** You will get sedatives which can greatly affect your judgment. You must have an adult to drive you home or your exam will be cancelled. You may take a cab or bus home; however, you must have a responsible adult with you other than the driver.
4. Our staff will need to review your health history. This includes any past surgeries, allergies, and medicines that you take. Please bring this information with you.
5. If you use **CPAP or BiPAP** for sleep apnea, please bring it with you. We will need your machine to safely sedate you.

The Day of Your Exam (Push, Single,

Double-Balloon)

1. No solid food, milk, or dairy products until after your exam.
2. Do not take liquid antacids before your exam.
3. You may drink clear liquids until **4 hours** before you arrive (i.e. water, apple juice, soda[®], Gatorade[®]). You must stop for sedation to be given safely. **Avoid red liquids.**
4. Take your normal medicines with a small sip of clear liquid up to 1 hour before you arrive. **Do not take the pills and supplements mentioned under “Important to Know.”**
5. **Follow the arrival and registration instructions in your cover letter.** If you are delayed, call us at **(608) 263-8094** to let us know you are going to be late. If you are over 30 minutes late, we may have to cancel and reschedule. We do our best to stay on time.

The Exam (Push, Single or Double Balloon)

A nurse will review your health history and place an IV. The IV is used to give you medicines which make you feel drowsy and relaxed during your test. The test will take about 1-2 hours.

After the test you will go to our recovery room for about 1 hour. Your driver may sit with you there. Once awake, you may be given something to drink. Your doctor will

discuss results with you before you go home. You and the doctor who referred you will get a copy of the report.

After the Exam and for the Rest of the Day

- Do not return to work.
- Do not drive.
- Do not use hazardous machinery.
- Do not make any important decisions.
- Plan to rest.
- Do not drink alcohol.
- You should be able to resume normal daily tasks on the day after your exam.
- You may have a mild sore throat. Salt water gargles should help.
- You may have some gas pains from the air used to distend your stomach. If you belch, this will help.

How to Contact Us GI Procedure Center 7:00 AM to 5:00 PM **(608) 263-8094**

After clinic hours, this number will give you the paging operator. Ask for the Gastroenterology (GI) doctor on call. Leave your name and phone number with the area code. The doctor will call you back.

If you live out of the area, call **1-800-323-8942**. **If you are not able to keep your appointment**, let us know 5 days in advance.

Your health care team may have given you this information as part of your care. If so, please use it and call if you have any questions. If this information was not given to you as part of your care, please check with your doctor. This is not medical advice. This is not to be used for diagnosis or treatment of any medical condition. Because each person's health needs are different, you should talk with your doctor or others on your health care team when using this information. If you have an emergency, please call 911. Copyright ©12/2018. University of Wisconsin Hospitals and Clinics Authority. All rights reserved. Produced by the Department of Nursing. HF#8067