

Home Care After IV Cytosan

- For the next 24 hours, have your child urinate every 2 hours while awake and every 4 hours while sleeping.
- Give your child plenty of fluids for the next 24 hours.
- Ondansetron _____ mg can be given every 8 hours, as needed, for nausea.

Call the doctor if your child has any severe vomiting or visible blood in the urine.

- In 10-14 days, have your doctor order a lab test. The test is called CBD with differential and platelets. The test results should be faxed to: **608-833-3211**, Attention: Pediatric Renal.

Your next appointment for IV Cytosan will be _____.

If you have any questions, please call the Pediatric Specialties Clinic at **(608) 263-6420**.

After hours, or if you live out of the area, please call **1-800-323-8942**.

If you are a patient receiving care at UnityPoint – Meriter, Swedish American or a health system outside of UW Health, please use the phone numbers provided in your discharge instructions for any questions or concerns.

Your health care team may have given you this information as part of your care. If so, please use it and call if you have any questions. If this information was not given to you as part of your care, please check with your doctor. This is not medical advice. This is not to be used for diagnosis or treatment of any medical condition. Because each person's health needs are different, you should talk with your doctor or others on your health care team when using this information. If you have an emergency, please call 911. Copyright © 4/2022 University of Wisconsin Hospitals and Clinics Authority. All rights reserved. Produced by the Department of Nursing. HF#5345.