

Wart treatment

Aldara Imiquimod 5% cream (12 sachets)

Ensure that you also read the enclosed instruction leaflet.

If you are pregnant **do not** use this cream.

Apply the cream at night, 3 times a week (on alternate days) for up to 4 months. E.g. on a Monday, Wednesday and Friday.


Take care not to use more cream than you need as this can damage your skin. If you have broken skin you will need to take a few days break from treatment to let your skin heal.

People with small clusters of warts do not need to use a full sachet for each application, and may choose to use a paper clip to save the cream for their next application - meaning the treatment lasts longer.

Wash the cream off before having sex and be aware that this cream may weaken latex condoms (non-latex condoms are available).

If you have visible symptoms then it is a good idea to talk to your current partner and avoid any skin to skin contact with the infected area, until you have completed treatment.

SH:24



You can use the plastic clip to reseal the sachet between applications.

Learn more:

Read more about genital warts and how they are treated here: sh24.org.uk/sexual-health

SH:24

**Sexual health:
24 hours a day**

Things to consider

- this cream does not clear the infection from your body completely. This means the warts may come back in the future
- don't apply this cream to warts inside the penis, vagina, or rectum
- if you experience any irritation or soreness, stop using the cream and contact us immediately
- when using this treatment, 50% of people find their warts clear in 8 weeks and 75% in 12 weeks*
- we'll text you in 11 weeks to see how well you have responded to treatment and whether you require another course.

*See instruction leaflet for more information



Talk to us:

If you need help or have questions, text: **07860 041 233**