SH:24

Sexual health: 24 hours a day

You have told us that:

- you have reported any medications you are taking to the SH:24 clinical team
- you do not have any medical problems that may be aggravated by this treatment (liver, kidney, heart, myasthenia gravis)
- you have reported any allergy to medications, peanuts or soya to the SH:24 clinical team
- if you are pregnant or breastfeeding you have discussed this with an SH:24 clinician
- you do not have any symptoms of chlamydia that may need to be reviewed in clinic e.g. abdominal pain.

Chlamydia treatment Azithromycin (4 x 500mg tablets)



SH:24

Ensure that you also read the A4 medication leaflet.

Day 1, take 2 tablets Day 2, take 1 tablet Day 3, take 1 tablet.

Do not chew them.

If you vomit within 3 hours of taking this treatment, get in touch as you may need an additional dose.

If you still have symptoms 7 days after you and your partner(s) have been treated, go to your local sexual health clinic as you may need further testing and treatment.

It is important to tell your partners that you have an infection so that they can get treated too.

Read more here: sh24.org.uk/get-support

Things to consider

- treatment takes 7 days to work
- avoid reinfection by not having sex until 7 days after you and your sexual partner(s) have completed the full course of treatment
- re-test 2 months after you have taken this treatment
- you may experience some diarrhoea after taking these tablets, if this is severe and is troubling you - get in touch
- condoms are very effective at preventing chlamydia infection, you can get free condoms at sexual health clinics.

Learn more:

Read more about chlamydia and how it is treated here: sh24.org.uk/sexual-health