

---

## Reference

David Perpetuini<sup>1</sup>, Emanuele Francesco Russo<sup>2</sup>, Daniela Cardone<sup>1</sup>, Roberta Palmieri<sup>3</sup>, Andreas De Giacomo<sup>3</sup>, Raffaello Pellegrino<sup>4</sup>, Arcangelo Merla<sup>1</sup>, Rocco Salvatore Calabrò<sup>5</sup> and Serena Filoni<sup>2</sup>.

<sup>1</sup>Department of Engineering and Geology, University G. D'Annunzio of Chieti-Pescara, Pescara, Italy;

<sup>2</sup> Padre Pio Foundation and Rehabilitation Centers, San Giovanni Rotondo, Italy;

<sup>3</sup> Translational Biomedicine and Neuroscience Department (DiBrain), University of Bari "Aldo Moro", Bari, Italy;

<sup>4</sup> Department of Scientific Research, Campus Ludes, Off-Campus Semmelweis University, Lugano, Switzerland;

<sup>5</sup> IRCCS Centro Neurolesi "Bonino-Pulejo", Messina, Italy.

# Use and Effectiveness of Electrosuit in Neurological Disorders: A Systematic Review with Clinical Implications

Bioengineering 2023, 10(6), 680. <https://doi.org/10.3390/bioengineering10060680>. (open-access article)

---

## Products

### Exopulse Mollii Suit (EMS)

---

## Major Findings

With Exopulse Mollii Suit:

#### → Improved mobility

- Children with CP improved in mobility and gross motor functions (significant in 1 of 3 publications)
- Stroke and CP patients improved in time and number of steps in the 10m comfortable gait test (1 publication)

#### → Improved function

- Significant improvements in upper and lower extremity functions (Fugl-Meyer Assessment) in stroke patients (1 publication)
- trunk acceleration in the anterior-posterior direction of children with CP was altered towards that of healthy individuals (1 publication)

#### → Decreased spasticity

- Spasticity measured with Modified Ashworth Scale (MAS) and Modified Tardieu scale (MTS) decreased significantly after treatment in children with CP (in 1 of 4 publications)
- Neural component of resistance to passive stretch decreased significantly in the wrist flexors of affected hand in stroke patients (1 publication)

#### → Improved active and passive Range of Motion

in children with CP (pROM, significant in 1 of 3 publications)

#### → Decreased pain

in children with CP and in adults with different types of pain (significant in 2 of 5 publications)

#### → Improved activity

Children with CP improved in goal-attainment scaling for self-selected activities (1 publication)

	Cerebral Palsy	Cerebral Palsy + Stroke	Stroke	Pain diagnoses (Fibromyalgia, Parkinson and others)
<b>Mobility</b>	↑↑ GMFM (1 publication) ○ GMFM (1 publication) ○ gross motor function (1 publication)	↑ 10m comfortable gait test		
<b>Function</b>	↑↑ Nonlinear dynamics of trunk acceleration		↑↑ FM-UE, FM-LE	
<b>Spasticity</b>	↓↓ MAS (1 publication) ↓↓ MTS (1 publication) ↓ MTS (1 publication) ○ MTS (1 publication)	○ MAS (1 publication)	○ MAS (2 publications)	
<b>Range of Motion</b>	↑↑ (1 publication) ○ (2 publications)			
<b>Pain</b>	↓ pain (1 publication) ○ (2 publications)			↓↓ VAS (1 publication) ↓↓ NRS (1 publication)
<b>Participation</b>	↑↑ COPM			


↓↓↓,↑↑↑ highly statistically significant ( $p < 0,001$ ) Increase/Reduction; ↑↑, ↓↓ statistically significant ( $p < 0,05$ ) Increase/Reduction; ↑, ↓ statistical trend ( $0 < p < 0.1$ ) towards Increase/Reduction; ○ no changes; GMFM Gross Motor Function Measure; MAS Modified Asworth Scale; MTS Modified Tardieu Scale; ROM Range of Motion; pROM passive Range of Motion, VAS Visual Analogue Scale; NRS Numerical Rating Scale, COPM Canadian Occupational Performance Measure;

## Population

Subjects: 336 subjects<sup>i</sup> (273 adults, 55 children)  
 Etiology: Cerebral Palsy (n = 63), Cerebral Palsy or Stroke (n= 42), Stroke (n = 40), Fibromyalgia (n = 73), Parkinson (n = 29), other types of pain (n = 89)  
 Individual study designs: 9 studies (n=273) with before-and-after-design  
 2 studies (n=47) with inactive/placebo EMS as control  
 1 study (n=16) with parallel control group (n=8) with conventional therapy

## Study Design

Systematic literature review



A systematic review was performed in accordance with Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) statement (Rethlefsen 2021). The systematic literature search was conducted in the databases PubMed, MEDLINE, Web of Science, and Scopus for literature from the inception to February 2023. The search concentrated on the words “Mollii” and “neurological disorders” and/or “stroke” and/or “PCI/cerebral palsy” and/or “rehabilitation” in the article titles, abstracts, and keywords.

<sup>i</sup> Discrepancy of n=10 for stated number of patients in publication of Riachi et al. 2019

## Results

Body Functions			Activity				Participation	Environment	Others
Biomechanical	Medical (Pain / Injuries)	Other	Upper extremity function	Lower extremity function	Safety	Activity, Mobility, ADLs	Preference, Satisfaction, QoL	Health Economics	Technical aspects

Category	Outcomes	Pathology	Results	Sig.*	
Biomechanics	Nonlinear dynamics of trunk acceleration	CP	Temporal structure of trunk acceleration in the anterior-posterior direction was altered towards that of healthy individuals. <sup>ii</sup> (Decrease in Largest Lyapunov exponent ( $p=0.041$ ) and complexity Index ( $p=0.030$ ))	++ Raffalt 2022	
Medical	Pain	VAS	Fibromyalgia, Parkinson, and other types of pain	VAS-0: $6.5 \pm 1.24$ (before intervention) VAS-1: $3.46 \pm 1.4$ (immediately afterwards) VAS-24: $4.72 \pm 1.68$ (24h later) Significant drop in VAS-1 and VAS-24 ( $p < 0.001$ )	-- Riachi 2019
	Numerical or graphical rating	CP	Pain was reduced in children who reported pain at the beginning of the research <sup>iii</sup> No changes	- 0 Flodström 2022 Arkkukangas 2022 Hedin 2022	
		Fibromyalgia (n=1)	Numerical Rating Scale (NRS) was reduced <sup>iv</sup>	-- Rubio-Zarapiz 2023	
Spasticity	Modified Asworth Scale (MAS)	CP	degree of spasticity decreased after one month ( $p=0.007$ ), and after six months ( $p=0.011$ ).	-- Hedin 2022	
		CP + stroke	No changes	0 Ertzgaard 2018	
		Stroke	No changes	0 Palmcrantz 2020 Pennati 2021	
	Modified Tardieu Scale	CP	Decreased after one month ( $p=0.030$ ) Decreased after three ( $p=0.392$ ) and six months ( $p=0.426$ ) No changes	-- Hedin 2022 - Hedin 2022 0 Bakaniene 2018	
	Muscle tone	CP	No changes	0 Flodström 2022	

Category	Outcomes	Pathology	Results	Sig.*
	Passive and Active Range of Motion (pROM, ROM)	CP	After one, three, and six months of treatment, a significant number of improved muscles were observed for pROM (p=0.000, p=0.001, p=0.014)	++ Hedin 2022
			No changes	0 Bakaniene 2018 Flodström 2022
		Stroke	No changes	0 Pennati 2021
	NeuroFlexor surface EMG	Stroke	No changes	0 Pennati 2021
	NeuroFlexor® Neural Component (NC)	Stroke	NC decreased in wrist flexors of affected hand (p=0.023) <sup>v</sup>	-- Palmcrantz 2020
Oxygenation	Hands temperature	Fibromyalgia (n=1)	Decrease in hands temperature <sup>iv</sup>	-- Rubio-Zarapiz 2023
	Muscle oxygen saturation SmO <sub>2</sub>	Fibromyalgia (n=1)	Increased <sup>iv</sup>	++ Rubio-Zarapiz 2023
	Total haemoglobin THb	Fibromyalgia (n=1)	No changes	0 Rubio-Zarapiz 2023
	Oxygenated haemoglobin O <sub>2</sub> Hb	Fibromyalgia (n=1)	Increased <sup>iv</sup>	++ Rubio-Zarapiz 2023
	Deoxygenated haemoglobin HHb	Fibromyalgia (n=1)	Decreased <sup>iv</sup>	-- Rubio-Zarapiz 2023
	Forced expiratory volume in 1s (FEV1)	Fibromyalgia (n=1)	Decreased <sup>iv</sup>	-- Rubio-Zarapiz 2023
	Forced expiratory volume in 6s (FEV6)	Fibromyalgia (n=1)	Increased <sup>iv</sup>	++ Rubio-Zarapiz 2023
	FEV1/FEV6	Fibromyalgia (n=1)	Decreased <sup>iv</sup>	-- Rubio-Zarapiz 2023
Vegetative system (relaxation)	Sleep	CP	No changes	0 Arkkukangas 2022 Hedin 2022
	Bowel function	CP	No changes	0 Hedin
	Cortical arousal	Fibromyalgia (n=1)	No changes	0 Rubio-Zarapiz 2023
	Saliva flux	Fibromyalgia (n=1)	Increased <sup>iv</sup>	++ (sic) Rubio-Zarapiz 2023
	Salivary proteins	Fibromyalgia (n=1)	Decreased <sup>iv</sup>	-- Rubio-Zarapiz 2023
Upper extremity function	Action Reach Arm Test (ARAT)	Stroke	Increased <sup>vi</sup>	++ (sic) Palmcrantz 2020
		CP + Stroke	No changes	0 Ertzgard 2018

Category	Outcomes	Pathology	Results	Sig.*
	Fugl-Meyer index - Upper Extremity total score	Stroke	Increased (p=0.000) <sup>vi</sup>	++ Palmcrantz 2020
	Level of Sitting Scale (LSS)	CP	No changes	0 Arkkukangas 2022
	Box and Block test	CP	No changes	0 Arkkukangas 2022
	Grip strength	Stroke	No changes	0 Palmcrantz 2020
		Fibromyalgia (n=1)	Improved <sup>iv</sup>	++ (sic) Rubio-Zarapiz 2023
	Wolf Motor function test (WMFT) – first two tasks	CP + Stroke	No changes	0 Ertzgaard 2018
Lower extremity function	Fugl-Meyer index - Lower Extremity total score	Stroke	Increased (p=0.003) <sup>vi</sup>	++ Palmcrantz 2020
	10m walk test	CP + Stroke	Comfortable gait test, time and number of steps improved <sup>vi</sup>	+ Ertzgaard 2018
		Stroke	No changes	0 Palmcrantz 2020
	6 min walk test	Stroke	No changes	0 Palmcrantz 2020
	10m up and go test	Fibromyalgia (n=1)	Improved <sup>iv</sup>	++ (sic) Rubio-Zarapiz 2023
	One leg balance	Fibromyalgia (n=1)	Improved <sup>iv</sup>	++ (sic) Rubio-Zarapiz 2023
	Chair stand test	Fibromyalgia (n=1)	Improved <sup>iv</sup>	++ (sic) Rubio-Zarapiz 2023
Safety	TUG [s]	CP	No changes	0 Bakaniene 2018 Arkkukangas 2022
		CP + Stroke	No changes	0 Ertzgaard 2018
	Berg Balance Scale (BBS)	Stroke	No changes	0 Palmcrantz 2020
Activity	Goal Attainment Scale (GAS)	CP + Stroke	Improved <sup>vi</sup>	+ Ertzgaard 2018
	Canadian occupational performance measure (COPM)	CP	All participants (n=6) showed improvements in total score of COPM, three of them showed significant clinical improvements <sup>iii</sup>	++ Flodström 2022
Mobility	Gross Motor Function Measure (GMFM)	CP	Increased <sup>v</sup>	++ Bakaniene 2018
			No changes	0 Hedin 2022

Category	Outcomes	Pathology	Results	Sig.*
	Gross Motor Function	CP	No changes	0 Flodström 2022
Preference, Satisfaction, Quality of Life (QoL)	Stroke Impact Scale	Stroke	No changes	0 Palmcrantz 2020
	Experiences with the suit (impact on body, self, activities, participation in ADLs)	CP	Children and parents saw improvements in children's physical and mental health after the use of the suit.	n.a. Nordstrom 2021
	Interest in continuing using Mollii, Motivation, Usability, Support	CP + Stroke	Responder participants felt hopeful, motivated when experiencing a treatment effect, and disappointed when not.	n.a. Jonasson 2022

**ADLs:** Activities of Daily Life; **ARAT:** Action Reach Arm Test; **BBS:** Berg Balance Scale; **CGT:** Comfortable gait test; **COPM:** Canadian occupational performance measure; **EMG:** Electromyography; **FEV:** Forced expiratory volume; **FEV1:** Forced expiratory volume in 1s; **FEV6:** Forced expiratory volume in 6s; **FGT:** Fast gait test; **FM:** Fugl-Meyer index; **GAS:** Goal Attainment Scale; **GMFM:** Gross Motor Function Measure; **LSS:** Level of sitting scale; **MAS:** Modified Asworth Scale; **MTS:** Modified Tardieu Scale, **NC:** Neural component; **PPT:** Pressure Pain Threshold; **pROM:** passive Range of Motion; **ROM:** active Range of Motion; **QoL:** Quality of Life; **TUG:** Timed-up and go-Test; **VAS:** Visual Analogue Scale; **WMFT:** Wolf motor function test

\* no difference (0), positive trend (+), negative trend (-), significant (++/--), not applicable (n.a.)

<sup>ii</sup> after 24 weeks of intervention; <sup>iii</sup> after 3 months of intervention; <sup>iv</sup> after 60 minutes of intervention; <sup>v</sup> after 3 weeks of intervention; <sup>vi</sup> after 6 weeks of intervention;

## Author's Conclusion

"The EMS is a full-body garment with built-in electrodes that deliver electrical stimulation to the muscles, with the goal of reducing spasticity and increasing flexibility and range of motion. This review investigated the benefits of the employment of the EMS for rehabilitation purposes for CP, stroke, and other neurological disease patients. Although further studies should be performed to test the effectiveness of this device, the literature reviewed demonstrated the potentiality of the EMS in clinical practice to improve the motor functions of neurological patients." (Perpetuini et al. 2021)

© 2023, Otto Bock HealthCare Products GmbH ("Otto Bock"), All Rights Reserved. This article contains copyrighted material. Wherever possible we give full recognition to the authors. We believe this constitutes a 'fair use' of any such copyrighted material according to Title 17 U.S.C. Section 107 of US Copyright Law. If you wish to use copyrighted material from this site for purposes of your own that go beyond 'fair use', you must obtain permission from the copyright owner. All trademarks, copyrights, or other intellectual property used or referenced herein are the property of their respective owners. The information presented here is in summary form only and intended to provide broad knowledge of products offered. You should consult your physician before purchasing any product(s). Otto Bock disclaims any liability related from medical decisions made based on this article summary.

## References

- Arkkukangas M, Hedberg Graff J, Denison E. Evaluation of the Electro-Dress Mollii® to Affect Spasticity and Motor Function in Children with Cerebral Palsy: Seven Experimental Single-Case Studies with an ABAB Design. *Cogent. Eng.* 2022;9:2064587.
- Bakaniene I, Urbonaviciene G, Janaviciute K, Prasauskiene A. Effects of the Interventions Method on Gross Motor Function in Children with Spastic Cerebral Palsy. *Neurol. I Neurochir. Pol.* 2018;52:581-586.
- Ertzgaard P, Alwin J, Sorbo A, Lindgren M, Sandjo L. Evaluation of a Self-Administered Transcutaneous Electrical Stimulation Concept for the Treatment of Spasticity: A Randomized Placebo-Controlled Trial. *Eur. J. Phys. Rehabil. Med.* 2018;54:507-517.

- Flodström C, Viklund Axelsson SA, Nordström BA. Pilot Study of the Impact of the Electro-Suie Mollii® on Body Functions, Activity, and Participation in Children with Cerebral Palsy. *Assist. Technol.* 2022;34:411-417.
- Hedin H, Wong C, Sjärdén A. The Effects of Using an Electrodress (Mollii®) to Reduce Spasticity and Enhance Functioning in Children with Cerebral Palsy: A Pilot Study. *Eur. J. Physiother.* 2022;24:134-143.
- Jonasson LL, Sörbo A, Ertzgaard P, Sandsjö L. Patients' experiences of self-administered electrotherapy for spasticity in stroke and cerebral palsy: A qualitative study. *J. Rehabil. Med.* 2022;54:1131.
- Nordstrom B, Prellwitz M. A Pilot Study of Children and Parents Experiences of the Use of a New Assistive Device, the Elctor Suit Mollii. *Assist. Technol.* 2021;33,238-245.
- Palmcrantz S, Pennati GV, Bergling H, Borg J. Feasibility and Potential Effects of Using the Electro-Dress Mollii on Spasticity and Functioning in Chronic Stroke. *J. Neuroeng. Rehabil.* 2020;17:1-10.
- Pennati GV, Bergling H, Carment L, Borg J, Lindberg PG, Palmcrantz S. Effects of 60 Min Electrostimulation with the EXOPULSE Mollii Suit on Objective Signs of Spasticity. *Front. Neurol.* 2021;12:706610.
- Raffalt PC, Bencke J, Mortensen K, Torabi TP, Wong , Speedtsberg MB. Elector-Suit Treatment of Children with Unilateral Cerebral Palsy Alters Nonlinear Dynamics of Walking. *Clin. Biomech.* 2022;98:105714.
- Rethlefsen ML, Kirtley S, Waffenschmidt S, Ayala AP, Moher D, Page MJ, Koeffel JB. PRISMA-S: An Extension to the PRISMA Statement for Reporting Literature Searches in Systematic Reviews. *Syst. Rev.* 2021;10:1-19.
- Riachi N, Khazen G, Ahdab R, Jörgen S. Pain Reducing Properties of the Mollii Suit on Adults with Chronic Pain Syndromes. *J. Neurol. Sci.* 2019;405:138-139.
- Rubio-Zarapiz A, Apolo-Arenas MD, Clemente-Suárez VJ, Costa AR, Pardo-Caballero D, Parraca JA. Acute Effects of a Session with The EXOPULSE Mollii Suit in a Fibromyalgia Patient: A Case Report. *Int. J. Environ. Res. Public Health* 2023;20:2209.