

PEEL HERE



Ultracare[®] ORAL ANESTHETIC GEL Walterberry[®]

1.01 FL OZ (30 mL) REF/UP 301

ULTRADENT
PRODUCTS, INC.

Made in USA | 33324.19 062518



Drug Facts

Active ingredient

Benzocaine 20% w/v

Purpose

Oral Anesthetic

Uses For the temporary relief of occasional minor irritation and pain associated with • minor dental procedures • sore mouth and throat • minor injury of the gums • canker sores • minor irritation of the mouth or gums caused by dentures or orthodontic appliances.

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. ➤

Drug Facts (continued)

This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: • pale, gray, or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy.

For external use only

Allergy alert: Do not use if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics, PABA compounds, or sunscreen. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsens, see your dentist or doctor promptly. ➤

Drug Facts (continued)

Do not use • if taking sulfonamides • for teething • in children under 2 years of age

When using this product: avoid contact with eyes.

Keep out of reach of children: in case of overdose, get medical help or contact a Poison Control Center right away. Do not exceed recommended dosage.

Directions

Adults and children 2 years of age and older

Apply a thin film to affected area. Allow to remain in place at least 1 minute and then spit out. Use up to 4 times daily or as directed by your dentist or doctor.

Drug Facts (continued)

Children between 2 and 12 years of age | Should be supervised in the use of this product.

Children under 2 years of age | Do not use.

Other information • Phenylketonurics: Contains Phenylalanine 3 mg per gram • Do not use if tamper-evident seal is broken • Store at room temperature

Inactive ingredients aspartame, D&C red no. 28, ethyl alcohol, FD&C red no. 40, glycerin, natural and artificial flavors, polyethylene glycol, sodium saccharin

Questions or comments? 800.552.5512



For Professional Use Only

Manufactured by Ultradent Products, Inc.
505 West Ultradent Drive (10200 South),
South Jordan, UT 84095

© 2018 Ultradent Products, Inc. All Rights Reserved. ultradent.com