



National Childhood Vaccination Program in Iceland as of October 2025

Age:	Vaccination against:
During pregnancy	Pertussis (<i>Boostrix</i> , given to a mother who is approx. 28 weeks pregnant)
0–6 months: RSV immunization (nirsevimab) at birth from October to March or in the all	6 weeks Rotavirus (oral drops; <i>RotaTeq</i>)
	3 months Rotavirus (oral drops; <i>RotaTeq</i>) DTaP-Hib-Polio (<i>Pentavac</i>) and PCV (<i>Vaxneuvance</i>)
	5 months Rotavirus (oral drops; <i>RotaTeq</i>) DTaP-Hib-Polio (<i>Pentavac</i>) and PCV (<i>Vaxneuvance</i>)
12 months	DTaP-Hib-Polio (<i>Pentavac</i>) PCV (<i>Vaxneuvance</i>) and menACWY (<i>MenQuadfi</i>)
18 months	MMR and VZV (<i>ProQuad</i>)
2,5 years	VZV (<i>Varilrix</i>)
4 years	DTaP (<i>Boostrix</i>)
12 years	MMR (<i>M-M-RVAXPRO</i>) HPV (<i>Gardasil 9</i> ; all genders)
14 years	DTaP-Polio (<i>Repevax</i>)

6–59 months:
Seasonal influenza vaccination (*Vaxigrip*) in the fall