
















Acetaminophen pediatric dosing directions

Acetaminophen	If possible, use weight to dose; otherwise use age. Single Oral Dose†							
Child's weight & age	Under 12 lbs Under 4 mos	12-17 lbs 4-11 mos	18-23 lbs 12-23 mos	24-35 lbs 2-3 yrs	36-47 lbs 4-5 yrs	48-59 lbs 6-8 yrs	60-71 lbs 9-10 yrs	72-95 lbs 11 yrs
Infants' Acetaminophen Concentrated Drops <i>for example</i> Infants' TYLENOL® Drops 80 mg per 1 mL	Talk to a doctor	 1 mL	 1.5 mL	—	—	—	—	—
Children's Acetaminophen Suspension Liquid <i>for example</i> Children's TYLENOL® Liquid 160 mg per 5 mL	← Use Infants' TYLENOL® Concentrated Drops as directed on the label or by a doctor (for infants' < 4 months of age) →			 5 mL (1 tsp)	 7.5 mL (1.5 tsp)	 10 mL (2 tsp)	 12.5 mL (2.5 tsp)	 15 mL (3 tsp)
Children's Acetaminophen Tablets <i>for example</i> Children's TYLENOL® Chewables 160 mg	← Use Infants' TYLENOL® Concentrated Drops as directed on the label or by a doctor (for infants' < 4 months of age) →			 1 Tablet	 1.5 Tablets	 2 Tablets	 2.5 Tablets	 3 Tablets
Children's Acetaminophen Powder <i>for example</i> Children's TYLENOL® Easy Dissolve 160 mg	← Use Infants' TYLENOL® Concentrated Drops as directed on the label or by a doctor (for infants' < 4 months of age) →			Use Children's TYLENOL® Liquid or Children's TYLENOL® Chewables as directed on the label		 2 Packs	 2 Packs	 3 Packs

Important instructions

Single dose may be repeated every 4 to 6 hours, as needed

Do **NOT** give more than 5 doses per day

Do **NOT** use with any other product containing Acetaminophen

mL = milliliter mos = months of age
 mg = milligram yrs = years of age
 lbs = pounds tsp = teaspoonful

What you can do













- Always read and follow the medicine label.
- Find the dose for your child's **current** weight. If you are not sure of your child's weight, use age. As children grow, their doses change over time.
- Use only the dosing device that came with the medicine (syringe or dosing cup).
- Wait the full amount of time between doses. If you give another dose too soon, it could harm your child.

See [ibuprofen dosing directions](#) ►

Find tips, quizzes, and videos at [GetReliefResponsibly.ca](https://www.getreliefresponsibly.ca)

Get Relief Responsibly®

Ibuprofen pediatric dosing directions

Ibuprofen	If possible, use weight to dose; otherwise use age. Single Oral Dose ^s							
	Under 12 lbs Under 4 mos	12-17 lbs 4-11 mos	18-23 lbs 12-23 mos	24-35 lbs 2-3 yrs	36-47 lbs 4-5 yrs	48-59 lbs 6-8 yrs	60-71 lbs 9-10 yrs	72-95 lbs 11 yrs
Infants' Ibuprofen Concentrated Drops for example Infants' MOTRIN® Concentrated Drops 40 mg per 1 mL	Talk to a doctor	 1.25 mL	 1.875 mL	—	—	—	—	—
Children's Ibuprofen Suspension Liquid for example Children's MOTRIN® 100 mg per 5 mL	← Use Infants' MOTRIN® Concentrated Drops as directed on the label or by a doctor (for infants' < 4 months of age) →			 5 mL (1 tsp)	 7.5 mL (1.5 tsp)	 10 mL (2 tsp)	 12.5 mL (2.5 tsp)	 15 mL (3 tsp)
Children's Ibuprofen Chewable Tablets for example Junior Strength Children's ADVIL® Chewables 100 mg	← Talk to a doctor →			 1 Tablet	 1.5 Tablets	 2 Tablets	 2.5 Tablets	 3 Tablets

Important instructions

Single dose may be repeated every 6 to 8 hours, as needed

Do NOT give more than 4 doses per day

Do NOT use with any other product containing Ibuprofen

mL = milliliter mos = months of age
 mg = milligram yrs = years of age
 lbs = pounds tsp = teaspoonful

What you can do

- Always read and follow the medicine label.
- Please consult product labelling or company websites for information regarding non-Johnson & Johnson manufactured products (e.g. Advil® (GSK/Haleon), including the most up-to-date information regarding product dosing.
- Find the dose for your child's **current weight**. If you are not sure of your child's weight, use age. As children grow, their doses change over time.
- Use only the dosing device that came with the medicine (syringe or dosing cup).
- Wait the full amount of time between doses. If you give another dose too soon, it could harm your child.

*nonsteroidal anti-inflammatory drug

[See acetaminophen dosing directions](#) ▶

Find tips, quizzes, and videos at [GetReliefResponsibly.ca](https://www.getreliefresponsibly.ca)

Get Relief Responsibly®