













# Acetaminophen pediatric dosing directions

Acetaminophen	If possible, use weight to dose; otherwise use age. Single Oral Dose†							
	Under 12 lbs Under 4 mos	12-17 lbs 4-11 mos	18-23 lbs 12-23 mos	24-35 lbs 2-3 yrs	36-47 lbs 4-5 yrs	48-59 lbs 6-8 yrs	60-71 lbs 9-10 yrs	72-95 lbs 11 yrs
<b>Infants' Acetaminophen Concentrated Drops</b> <i>for example</i> <b>Infants' TYLENOL® Drops</b> 80 mg per 1 mL	Talk to a doctor	 1 mL	 1.5 mL	—	—	—	—	—
<b>Children's Acetaminophen Suspension Liquid</b> <i>for example</i> <b>Children's TYLENOL® Liquid</b> 160 mg per 5 mL	← Use Infants' TYLENOL® Concentrated Drops as directed on the label or by a doctor (for infants' < 4 months of age) →			 5 mL (1 tsp)	 7.5 mL (1.5 tsp)	 10 mL (2 tsp)	 12.5 mL (2.5 tsp)	 15 mL (3 tsp)
<b>Children's Acetaminophen Tablets</b> <i>for example</i> <b>Children's TYLENOL® FASTMELTS® 160 mg</b>	← Use Infants' TYLENOL® Concentrated Drops as directed on the label or by a doctor (for infants' < 4 months of age) →			 1 Tablet	 1.5 Tablets	 2 Tablets	 2.5 Tablets	 3 Tablets

## Important instructions

Single dose may be repeated every 4 to 6 hours, as needed

Do **NOT** give more than 5 doses per day

Do **NOT** use with any other product containing Acetaminophen

### † For Infants' and Children's TYLENOL® products

- Single dose may be repeated every 4 to 6 hours, as needed
- Do **NOT** give more than 5 doses per day
- Do **NOT** use with any other product containing acetaminophen

mL = milliliter    mos = months of age  
 mg = milligram    yrs = years of age  
 lbs = pounds    tsp = teaspoonful

## What you can do








- Always read and follow the medicine label.
- Find the dose for your child's **current weight**. If you are not sure of your child's weight, use age. As children grow, their doses change over time.
- Use **only the dosing device that came with the medicine** (syringe or dosing cup).
- Wait the full amount of time between doses. If you give another dose too soon, it could harm your child.

See [ibuprofen dosing directions](#) ►

Find tips, quizzes, and videos at [GetReliefResponsibly.ca](https://www.getreliefresponsibly.ca)

Get Relief Responsibly®

# Ibuprofen pediatric dosing directions

Ibuprofen	If possible, use weight to dose; otherwise use age. Single Oral Dose <sup>§</sup>							
	Under 12 lbs Under 4 mos	12-17 lbs 4-11 mos	18-23 lbs 12-23 mos	24-35 lbs 2-3 yrs	36-47 lbs 4-5 yrs	48-59 lbs 6-8 yrs	60-71 lbs 9-10 yrs	72-95 lbs 11 yrs
<b>Child's weight &amp; age</b> <b>Infants' Ibuprofen Concentrated Drops</b> <i>for example</i> <b>Infants' MOTRIN® Concentrated Drops</b> <b>40 mg per 1 mL</b>	Talk to a doctor	 1.25 mL	 1.875 mL	—	—	—	—	—
<b>Children's Ibuprofen Suspension Liquid</b> <i>for example</i> <b>Children's MOTRIN® Oral Suspension</b> <b>100 mg per 5 mL</b>	Use Infants' MOTRIN® Concentrated Drops as directed on the label or by a doctor (for infants' < 4 months of age)			 5 mL (1 tsp)	 7.5 mL (1.5 tsp)	 10 mL (2 tsp)	 12.5 mL (2.5 tsp)	 15 mL (3 tsp)

**Important instructions**

Single dose may be repeated every 6 to 8 hours, as needed

Do **NOT** give more than 4 doses per day

Do **NOT** use with any other product containing ibuprofen

## § For Infants' and Children's MOTRIN® products

- Single dose may be repeated every 6 to 8 hours, as needed
- Do **NOT** give more than 4 doses per day
- Do **NOT** use with any other product containing ibuprofen

mL = milliliter    mos = months of age  
 mg = milligram    yrs = years of age  
 lbs = pounds    tsp = teaspoonful

## What you can do

- Always read and follow the medicine label.
- Find the dose for your child's **current** weight. If you are not sure of your child's weight, use age. As children grow, their doses change over time.
- Use only the dosing device that came with the medicine (syringe or dosing cup).
- Wait the full amount of time between doses. If you give another dose too soon, it could harm your child.

\*nonsteroidal anti-inflammatory drug

[See acetaminophen dosing directions](#) ▶

Find tips, quizzes, and videos at [GetReliefResponsibly.ca](https://www.getreliefresponsibly.ca)

Get Relief Responsibly®