

DEMONSTRATED PROTOCOL

Nuclei Isolation for Single Cell ATAC Sequencing

Overview

This protocol outlines how to isolate, wash, and count nuclei suspensions for use with the Chromium Single Cell ATAC protocol. Cryopreserved primary cells (PBMCs) and cell lines (GM12878 cells; EL4 cells) were used to develop this protocol. PBMCs were cryopreserved in IMDM + 40% FBS + 15% DMSO. Cell lines were cryopreserved in RPMI + 15% FBS + 5% DMSO. Optimization of some protocol steps (e.g. lysis time, centrifugation speed/time and filtration steps) may be needed based on cell type.

! The recommended buffer compositions, final nuclei suspension concentration, and the wash step guidelines presented in this protocol for nuclei sample preparation are critical for optimal Chromium Single Cell ATAC assay performance. Failure to adhere to these guidelines may result in compromised microfluidics chip operation.

Additional Guidance

Consult Demonstrated Protocol Cell Preparation Guide (Document CG000053) for Tips & Best Practices.

Cells carry potentially hazardous pathogens. Follow material supplier recommendations and local laboratory procedures and regulations for the safe handling, storage and disposal of biological materials.

Cell Sourcing

Cell Type	Species	Supplier
GM12878	Human	Coriell Institute
EL4	Mouse	ATCC
Normal Peripheral Blood MNC (PBMC)	Human	AllCells

Preparation – Buffers

Diluted Nuclei Buffer Maintain at 4°C	Stock	Final	1 ml
Nuclei Buffer (10x Genomics, PN-2000153*)	20X	1X	50 µl
Nuclease-free Water	-	-	950 µl

Wash Buffer	Stock	Final	10 ml
Prepare fresh, maintain at 4°C			
Tris-HCl (pH 7.4)	1M	10 mM	100 µl
NaCl	5M	10 mM	20 µl
MgCl ₂	1M	3 mM	30 µl
BSA	10%	1%	1 ml
Tween-20	10%	0.1%	100 µl
Nuclease-free Water	-	-	8.75 ml

Lysis Buffer	Stock	Final	5 ml
Prepare fresh, maintain at 4°C			
Tris-HCl (pH 7.4)	1 M	10 mM	50 µl
NaCl	5 M	10 mM	10 µl
MgCl ₂	1 M	3 mM	15 µl
Tween-20	10%	0.1%	50 µl
Nonidet P40 Substitute (if using Sigma (74385) 100% solution, prepare a 10% stock)	10%	0.1%	50 µl
Digitonin (incubate at 65°C to dissolve precipitate before use)	5%	0.01%	10 µl
BSA	10%	1%	500 µl
Nuclease-free Water	-	-	4.315 ml

Additional Buffers

RPMI + 10% FBS (maintain at 4°C, pre-warm at 37°C before use)

PBS + 0.04% BSA (maintain at 4°C)


Specific Reagents & Consumables

Vendor	Item	Part Number
10x Genomics	Nuclei Buffer	2000153*
Thermo Fisher Scientific	Digitonin Tubes, 0.2 ml, flat cap tube**	BN2006 AB0620
Fisher Scientific	Sorvall Microtube Adapters**	76003750
Millipore-Sigma	Trizma Hydrochloride Solution, pH 7.4 Sodium Chloride Solution, 5M Magnesium Chloride Solution, 1M Nonidet P40 Substitute	T2194 59222C M1028 74385
Miltenyi Biotec	MACS BSA Stock Solution	130-091-376
Bel-Art	Flowmi Cell Strainer, 40 µm	H13680-0040

*Included in the Chromium Single Cell ATAC Library Kit

**ONLY for Low Cell Input Nuclei Isolation protocol

Protocol Overview



Protocol

1. Thaw Cells (if using frozen cells)

For cell lines (used for GM12878 and EL4 cells):

- Remove cryovials from storage, thaw in the water bath at **37°C** for **1–2 min**. Remove from the water bath when a tiny ice crystal remains.
- Pipette mix the cells and transfer to a 15-ml conical tube containing **10 ml** pre-warmed media (RPMI + 10% FBS).
- Centrifuge at **300 rcf** for **5 min**.
- Remove the supernatant without disrupting the cell pellet and resuspend in **1 ml** PBS + 0.04% BSA. Transfer to a 2-ml microcentrifuge tube. Rinse the 15-ml tube with **0.5 ml** PBS + 0.04% BSA and transfer the rinse to the 2-ml tube containing the cells.
- Centrifuge cells at **300 rcf** for **5 min**.
- Remove the supernatant without disrupting the cell pellet and resuspend in **1 ml** PBS + 0.04% BSA.
- Pass cell suspension through a **40 µm** Flowmi Cell Strainer.
- Determine the cell concentration using a Countess II FL Automated Cell Counter (see Appendix) or a hemocytometer.
- Proceed to Nuclei Isolation (step 2).
If cell count is <100,000, nuclei may be isolated using the Low Cell Input Nuclei Isolation protocol (see Appendix).

For primary cells/fragile cells (used for PBMCs):

- Remove cryovial from storage, thaw in the water bath at **37°C** for **2-3 min**. Remove from the water bath when a tiny ice crystal remains.
- Transfer the thawed cells to a 50-ml conical tube. Rinse the cryovial with **1 ml** pre-warmed media (RPMI + 10% FBS) and add the rinse drop-wise to the 50-ml conical tube while gently shaking the tube.
- Serially dilute cells with media a total of 5 times (including dilution at step b) by 1:1 volume additions with **~1 min** wait between additions (see Appendix). Add media (RPMI + 10% FBS) at a speed of **1 ml/3-5 sec**.
- Centrifuge at **300 rcf** for **5 min**.
- Remove most of the supernatant, leaving **~1 ml** and resuspend cell pellet in this volume.
- Add an additional **9 ml** media (**1 ml/3-5 sec**) to achieve a total volume of **~10 ml**.
- Centrifuge at **300 rcf** for **5 min**.
- Remove the supernatant without disrupting the cell pellet and resuspend in **1 ml** PBS + 0.04% BSA gently pipette mix 5x.
- Transfer to a 2-ml microcentrifuge tube. Rinse the 50-ml tube with **0.5 ml** PBS + 0.04% BSA and transfer the rinse to the 2-ml tube containing the cells. Resuspend by gently inverting the tube.
- Centrifuge cells at **300 rcf** for **5 min**.

- Remove the supernatant without disrupting the cell pellet and resuspend in **1 ml** PBS + 0.04% BSA.
- Pass cell suspension through a **40 µm** Flowmi Cell Strainer.
- Determine the cell concentration using a Countess II FL Automated Cell Counter (see Appendix) or a hemocytometer.
- Proceed to Nuclei Isolation (step 2).
If cell count is <100,000, nuclei may be isolated using the Low Cell Input Nuclei Isolation protocol (see Appendix).

2. Nuclei Isolation

If using fresh cells, perform 1-2 washes with PBS + 0.04% BSA and determine cell count before proceeding to step 2a. Nuclei may be isolated from 100,000-1,000,000 cells using this protocol.

- Add 100,000-1,000,000 cells to a 2-ml microcentrifuge tube. Centrifuge at **300 rcf** for **5 min** at **4°C**.
- Remove ALL the supernatant without disrupting the cell pellet.
- Add **100 µl** chilled Lysis Buffer. Pipette mix 10x.
- Incubate for **3-5 min*** on ice.
*Cryopreserved PBMCs were incubated for **3 min**
*Cryopreserved cell lines were incubated for **5 min**



Optimize incubation time based on cell type. Sub-optimal or prolonged lysis times can both alter assay performance. Assess lysis efficacy via the Countess II FL Automated Cell Counter/microscopy (see Results).


- Add **1 ml** chilled Wash Buffer to the lysed cells. Pipette mix 5x.
- Centrifuge at **500 rcf** for **5 min** at **4°C**.
- Remove the supernatant without disrupting the nuclei pellet.
- Based on cell concentration step 2a and assuming **~50%** nuclei loss during cell lysis, resuspend in chilled Diluted Nuclei Buffer. See Nuclei Stock Concentration Table and Example Calculation in Appendix. Maintain on ice.



The resuspension in Diluted Nuclei Buffer is critical for optimal Single Cell ATAC assay performance. The composition of the Tris-based Diluted Nuclei Buffer, including Magnesium concentration, has been optimized for the Transposition and Barcoding steps in the Single Cell ATAC protocol. Suspension of nuclei in a different buffer may not be compatible.

- OPTIONAL** If cell debris and large clumps are observed, pass through a cell strainer. For low volume, use a **40 µm** Flowmi Cell Strainer to minimize volume loss.
- Determine the nuclei concentration using a Countess II FL Automated Cell Counter (see Appendix) or a hemocytometer.
- Proceed **immediately** to Chromium Single Cell ATAC Reagent Kits protocol (CG000168).


Results



Troubleshooting


Problem	Possible Solution
High fraction of non-viable cells in input material prior to starting nuclei isolation	Optimize cell thawing to enhance sample quality
	Reduce fraction of dead cells. Refer to Demonstrated Protocol Removal of Dead Cells from Single Cell Suspensions for Single Cell RNA Sequencing (Document CG000093)
	Sort cells using flow cytometry
High fraction of viable cells post cell lysis	Gently handle cell suspensions by following best practices and reduce cell processing times
	Incrementally increase the lysis time and monitor lysis efficacy microscopically

Trypan Blue Precipitate in the Countess II Slide




DO NOT use nuclei resuspended in 20X Nuclei Buffer. Repeat nuclei isolation and resuspend in Diluted Nuclei Buffer (1X).

Nuclei Quality - Representative Images (Panel A: recommended quality)



Appendix

Illustrative Overview of 1:1 Serial Dilutions



Nuclei Stock Concentration Table

Based on the Targeted Nuclei Recovery, prepare the nuclei suspension in Diluted Nuclei Buffer to achieve the corresponding Nuclei Stock concentrations.

Targeted Nuclei Recovery	Nuclei Stock Concentration (nuclei/ μl)
500	155-390
1,000	310-780
2,000	610-1,540
3,000	925-2,300
4,000	1,230-3,075
5,000	1,540-3,850
6,000	1,850-4,600
7,000	2,150-5,400
8,000	2,460-6,150
9,000	2,770-6,900
10,000	3,080-7,700

Example Calculation

Cell count at step 2a: **200,000**
 Estimated nuclei count at step 2h (~50% loss): **100,000**
 If targeting 5,000 Nuclei Recovery, nuclei pellet at step 2h may be resuspended in **30 μl** Diluted Nuclei Buffer for Nuclei Stock Concentration of 1,540-3,850 nuclei/ μl (see Table above)

Appendix

Nuclei Counting and Viability

Countess II FL Automated Cell Counter is recommended for determining nuclei concentrations. The optimal range of cell concentration for Cell Counter is 1,000-10,000 cells/ μl . Refer to manufacturer's instructions for details on operations.

- Vortex 0.4% trypan blue stain, centrifuge briefly and aliquot **10 μl** per tube.
- Pipette mix the nuclei suspension. Immediately add 10 μl nuclei suspension to 10 μl aliquot of 0.4% trypan blue stain. Gently pipette mix 10x.
- Transfer 10 μl trypan blue stained nuclei to a Countess II Cell Counting Slide chamber.
- Insert the slide into the Countess II FL Cell Counter, and determine the nuclei concentration and viability. <5% of input cells should be viable. Optimize focusing and light exposure.

Low Cell Input Nuclei Isolation

Nuclei may be isolated from 2,000-100,000 cells using this protocol. If cell count is <40,000, centrifuge cell suspension at **300 rcf for 5 min at 4°C** and resuspend the cell pellet in **50 μl PBS + 0.04% BSA**. Transfer **50 μl** cell suspension to a 0.2-ml tube. Proceed directly to **step c**.

- Centrifuge cell suspension at **300 rcf for 5 min at 4°C**. Remove supernatant and resuspend pellet in **PBS + 0.04% BSA** for 1,000 cells/ μl cell suspension.
- Add 2,000-40,000 cells to a 0.2-ml tube in a total volume of **50 μl PBS + 0.04% BSA**.

Approximately 25% of the cell input is expected to be recovered during Chromium Single Cell ATAC sequencing. Always determine nuclei counts after nuclei isolation.


Cell Input	Expected Nuclei Recovery (after cell lysis)	Expected Nuclei Recovery (ATAC sequencing)
40,000	16,000	10,000
20,000	8,000	5,000
10,000	4,000	2,500
4,000	1,600	1,000
2,000	800	500

- Centrifuge at **300 rcf for 5 min at 4°C**.
- Remove **45 μl** supernatant without touching the bottom of the tube to avoid dislodging the cell pellet.
- Add **45 μl** chilled Lysis Buffer. Gently pipette mix 3x.
- Incubate for **3-5 min*** on ice.
 - *Cryopreserved PBMCs were incubated for **3 min**
 - *Cryopreserved cell lines were incubated for **5 min**



Optimize incubation time based on cell type. Sub-optimal or prolonged lysis times can both alter assay performance. Assess lysis efficacy via the Countess II FL Automated Cell Counter/microscopy. See Results for optimal cell lysis.

- Add **50 μl** chilled Wash Buffer to the tube. **DO NOT** mix.
- Centrifuge at **500 rcf for 5 min at 4°C**.
- Remove **95 μl** supernatant without disrupting the nuclei pellet.
- Add **45 μl** chilled Diluted Nuclei Buffer to the pellet. **DO NOT** mix.
- Centrifuge at **500 rcf for 5 min at 4°C**.
- Remove the supernatant without touching the bottom of the tube to avoid dislodging the nuclei pellet.



The supernatant may be removed in two steps, first with a 100- μl pipette (set to 40 μl), followed by removal with a 10- μl pipette (set to 10 μl).

- Resuspend the nuclei pellet in **7 μl** chilled Diluted Nuclei Buffer (pellet may not be visible).



The use of the Diluted Nuclei Buffer for nuclei suspension is critical for optimal Single Cell ATAC assay performance. The composition of the Tris-based Diluted Nuclei Buffer, including Magnesium concentration, has been optimized for the Transposition and Barcoding steps in the Single Cell ATAC protocol. Suspension of nuclei in a different buffer may not be compatible with these protocol steps.

- Use **2 μl** nuclei suspension mixed with **8 μl** Diluted Nuclei Buffer and **10 μl** Trypan Blue to determine the cell concentration by a Countess II FL Automated Cell Counter (see Appendix) or a hemocytometer. A final nuclei concentration of 30 nuclei/ μl is needed for Targeted Nuclei Recovery of 500.
- Proceed **immediately** with **5 μl** nuclei suspension to Chromium Single Cell ATAC Reagent Kits protocol (CG000168).

© 2019 10x Genomics, Inc. (10x Genomics). All rights reserved. Duplication and/or reproduction of all or any portion of this document without the express written consent of 10x Genomics, is strictly forbidden. Nothing contained herein shall constitute any warranty, express or implied, as to the performance of any products described herein. Any and all warranties applicable to any products are set forth in the applicable terms and conditions of sale accompanying the purchase of such product. 10x Genomics provides no warranty and hereby disclaims any and all warranties as to the use of any third-party products or protocols described herein. The use of products described herein is subject to certain restrictions as set forth in the applicable terms and conditions of sale accompanying the purchase of such product. A non-exhaustive list of 10x Genomics' marks, many of which are registered in the United States and other countries can be viewed at: www.10xgenomics.com/trademarks. 10x Genomics may refer to the products or services offered by other companies by their brand name or company name solely for clarity, and does not claim any rights in those third-party marks or names. 10x Genomics products may be covered by one or more of the patents as indicated at: www.10xgenomics.com/patents. The use of products described herein is subject to 10x Genomics Terms and Conditions of Sale, available at www.10xgenomics.com/legal-notices, or such other terms that have been agreed to in writing between 10x Genomics and user. All products and services described herein are intended FOR RESEARCH USE ONLY and NOT FOR USE IN DIAGNOSTIC PROCEDURES.

The use of 10x Genomics products in practicing the methods set forth herein has not been validated by 10x Genomics, and such non-validated use is NOT COVERED BY 10x GENOMICS STANDARD WARRANTY, AND 10x GENOMICS HEREBY DISCLAIMS ANY AND ALL WARRANTIES FOR SUCH USE. Nothing in this document should be construed as altering, waiving or amending in any manner 10x Genomics terms and conditions of sale for the Chromium Controller or the Chromium Single Cell Controller, consumables or software, including without limitation such terms and conditions relating to certain use restrictions, limited license, warranty and limitation of liability, and nothing in this document shall be deemed to be Documentation, as that term is set forth in such terms and conditions of sale. Nothing in this document shall be construed as any representation by 10x Genomics that it currently or will at any time in the future offer or in any way support any application set forth herein.

Contact:
support@10xgenomics.com
 10x Genomics
 7068 Koll Center Parkway
 Suite 401
 Pleasanton, CA 94566 USA

