

Polymorphous Adenocarcinoma (PAC) Presenting As a Palatal Swelling- A Case Report

Riya Jain^{1*}, Suchitra Gosavi², Deepak Sethia³ and Akshay Trimukhe⁴

¹Department of Oral Pathology, Government dental college and hospital Nagpur, India

²Department of Oral Pathology, Government dental college and hospital Nagpur, India

³Department of Critical Care Medicine, Deenanath Mangeshkar hospital, India

⁴Department of Oral and Maxillofacial Pathology, India

Citation: Riya Jain, Suchitra Gosavi, Deepak Sethia, Akshay Trimukhe. Polymorphous Adenocarcinoma (PAC) Presenting As a Palatal Swelling- A Case Report. *Ann Case Rep Clin Stud*. 2023;2(1):1-11.

Received Date: 28 September, 2022; **Accepted Date:** 16 December, 2022; **Published Date:** 13 January, 2023

***Corresponding author:** Riya Jain. Department of Oral Pathology, Government dental college and hospital Nagpur, India

Copyright: © Riya Jain, Open Access 2023. This article, published in *Ann Case Rep Clin Stud (ACRCS)* (Attribution 4.0 International), as described by <http://creativecommons.org/licenses/by/4.0/>.

1. ABSTRACT

1.1. Background: Polymorphous adenocarcinoma is the second most common intraoral minor salivary gland tumor accounting for 26% of all carcinoma. It is characterized by varied architectural patterns with uniform cytologic picture.

1.2. Methods: We present a case of 45-year-old male who presented with a palatal swelling. neoplastic glandular epithelial cells showing variety of growth patterns and characteristic indian file pattern was also seen. Micro calcifications were also appreciated.

1.3. Results: All these features are suggestive of Polymorphous Adenocarcinoma (PAC). The patient received postoperative radiotherapy in view of close margins for a total dose of 50.4 Gy in 28 fractions. The patient is doing well 33 months post surgery without any clinical or radiological evidence of recurrence.

1.4. Conclusions: Palatal salivary gland malignancies pose a diagnostic dilemma for the pathologists. Any general practitioner should be skeptical about the palatal swelling and refer the patient to a specialist.

2. INTRODUCTION

Palate being rich in minor salivary glands creates a possibility of a wide array of malignancies and poses a diagnostic dilemma. [1] Hence clinicians should be skeptical when a patient presents with palatal swelling. Batsakis first described Polymorphous Adenocarcinoma (PAC) as Terminal Duct Carcinoma. [2] Later Freedman and Lumerman described it as Lobular carcinoma in 1983. [3] However, Evans and Batsakis suggested the term “Polymorphous Low Grade Adenocarcinoma” (PLGA). Recently, in 2017 WHO opted for the term “Polymorphous

Adenocarcinoma” (PAC). WHO in the recent classification includes “cribriform adenocarcinoma of minor salivary glands” (CAMSg) as a subtype of PAC. [4] Polymorphous adenocarcinoma is the second most common intraoral minor salivary gland tumor accounting for 26% of all carcinoma. [5] PAC shows a female predilection with a wide age range ranging from fifth to sixth decade of life. [6] Most commonly it presents as a painless mass on the palate. [5] PAC is characterized by varied architectural patterns with uniform cytologic picture. [7] On FNAC, cytological features may mimic Pleomorphic Adenoma (PA), Adenoid Cystic Carcinoma (ACC), mucoepidermoid carcinoma, epithelial myoepithelial carcinoma and papillary cystadenocarcinoma. [8] We present a case of 45-year-old male who presented with a palatal swelling.

3. CASE REPORT

A 45-year old male came to our department with the complaint of swelling over the left side of palate since 3 to 4 months. (Figure 1) The swelling increased in size drastically. He visited a private dentist for the same, where medications were prescribed for pain but no relief was seen. Thereafter, patient was referred to our department for further management. During the examination, a well demarked swelling was present over palatal region in 1st, 2nd, 3rd molar region on left side, Swelling was firm, non-tender in nature extending medially from mid palatal raphe to junction of horizontal process of palate and alveolar process. (Figure 1) Haematological and biochemical tests were within normal limits. Chest X-ray and ultrasound of abdomen and pelvis were normal. Cone Beam Computed Tomography (CBCT) was suggestive of a bulky left palatal lesion with is to hyperdense lesion. Fine needle aspiration cytology from the parotid swelling was suggestive of pleomorphic adenoma.

The patient underwent surgical excision of the tumor. Gross examination of the specimen revealed a grey white infiltrating tumor measuring 2.5 cm x 2 cm x 2 cm with focal grey brown areas. A histopathological examination was advised.

H & E stained section consists of parakeratinised stratified squamous surface epithelium which is separated at places from the lesional tissue by band of normal connective tissue. (Figure 2) The lesional tissue is partially encapsulated and consists of sheets of neoplastic glandular epithelial cells showing variety of growth pattern which includes cord like, duct like arrangement, cribriform pattern (Figure 3 and 4) peripheral part of section shows Indian file type of arrangement. (Figure 5)

These neoplastic glandular epithelial cells are round to oval in shape with scanty cytoplasm and pale staining vesicular nuclei. At one focus cells are arranged in a concentric pattern forming ‘targetoid’ appearance. (Figure 6)

At places, the cells are arranged in nests with intervening connective tissue septae resembling alveolar pattern. The cells are round to oval with granular eosinophilic cytoplasm and hyperchromatic nuclei. Clear cell differentiation is evident at places. (Figure 7)

The intervening connective tissue shows numerous areas of hyalinisation and few areas of mucous. (Figure 8 and 9)

At places areas of micro calcifications were seen. (Figure 10)

These neoplastic glandular epithelial cells have invaded the surrounding capsule and adjacent minor salivary gland tissue at places.

All these features are suggestive of Polymorphous Adenocarcinoma (PAC).

The patient received postoperative radiotherapy in view of close margins for a total dose of 50.4 Gy in 28 fractions.

The patient is doing well 33 months post surgery without any clinical or radiological evidence of recurrence.



Figure 1: Intraoral photograph of the patient showing palatal swelling on the left side of palate.




Figure2: Parakeratinised stratified squamous surface epithelium which is separated at places from the lesional tissue by band of normal connective tissue.




Figure3a: Duct like arrangement of neoplastic glandular cells (40x magnification).




Figure 3b: Opening up and channelization of tubules.




Figure 4: Cribriform arrangement resembling adenoid cystic carcinoma.




Figure 5: Single file/ Indian file arrangement of neoplastic cells.




Figure 6: Targetoid arrangement of cells in concentric layers.




Figure 7: Clear cells resembling mucoepidermoid carcinoma like areas.




Figure 8: Hyalinised stroma.




Figure 9: Mucoid stroma.




Figure10: Microcalcifications resembling psammoma bodies.

4. DISCUSSION

As early as 1983, Freedman and Lumermann published very first description of PAC when they termed it as lobular carcinoma to emphasize the similarity with single-file (Indian file) pattern seen in breast lobular carcinoma. [3] Simultaneously, Batsakis et al reported a series of similar tumors and proposed the term “terminal duct - adenocarcinoma” to denote their origin from terminal segment of salivary gland ducts. [2] It was in 1984 that Evans and Batsakis emphasized that this group of salivary gland tumors are characterized by bland uniform cytology but diverse architectural pattern and coined the term “Polymorphous Low Grade Adenocarcinoma” (PLGA). [8] However with experience it was understood that PLGA is not as indolent as was thought [9]. High grade transformation was also noted. [10] It was on these grounds that in 2017, WHO dropped out “Low Grade” and opted for the term “Polymorphous Adenocarcinoma” (PAC) and defined it as “a malignant epithelial tumor characterized by cytological uniformity, morphologic diversity and an infiltrative growth pattern”. [11]

PAC is the second most common salivary gland malignancy after Mucoepidermoid Carcinoma. [12] This tumor like any other salivary gland tumor affects women with F: M ratio of 2:1. [9] More than 90% PACs occur in more than 40 years of age with incidence rates between 40-79 years and mean age of diagnosis is 61.3 years affecting whites more than the people of color. [12] PAC is most commonly located on the posterior hard palate and soft palate. Buccal mucosa, labial mucosa may also be involved. [12,13] Upto 9% of PAC originate from major salivary gland predominantly parotid gland. [14] Most PACs present with approximately 2 cm size. [12] Few cases show pain, ulceration, bleeding or ill-fitting dentures. Our case was also in line with these findings except patient was a male. PAC has non-specific findings on imaging but can be an adjunct in assessing the local extent in the bone. [15] In our case Cone Beam Computed Tomography (CECT) was suggestive of a bulky left palatal lesion with iso to hyperdense lesion. Additional chest, abdominal radiographs are required to rule out metastasis [16,17] in our case both were normal.

For evaluation of salivary gland lesions, Fine needle aspiration cytology (FNAC) is very informative. [18] In our case the aspirate showed round basaloid cells and a diagnosis of pleomorphic adenoma was proposed with a differential diagnosis of Adenoid cystic carcinoma, Mucoepidermoid carcinoma. [19]

Diagnosis of PAC relies on incisional biopsy with margins of normal tissue. [20] Macroscopically, PAC is firm to solid with no capsule but circumscribed tumor mass lying just below the epithelium. The cut surface appears white or tan. [20] Tumor mass is adjacent to normal salivary gland lobules and flood the lamina propria just reaching the surface epithelium. [21] PAC is characterized by uniform appearance, round or polygonal shaped small to medium cells with indistinct boundaries. All dysplastic features like increased nucleo cytoplasmic N:C ration, vesicular nuclei with inconspicuous nucleoli are seen. These cytologic features contribute to the pale appearance of the tumor and was useful diagnostic criteria in the past. [5,9,21,22] Nuclear atypia and mitoses are not common.

PAC is famous for its varied architectural patterns ranging from solid islands & chords, tubules, pseudo-cribriform aggregates and indian-file pattern.

These pseudo-cribriform pattern which exhibits true lumina should be differentiated from classic cribriform pattern in Adenoid cystic carcinoma wherein they result from stromal cores trapped in tumor parenchyma and thus outlined

by basement membrane. [22] Targetoid pattern i.e. cells arranged concentrically around a nerve bundle is a common finding. [11,9]

Tyrosine rich crystals and microcalcifications can also be seen. [23] In our case also calcific bodies were seen.

For cytoskeleton and cytoplasmic filaments PAC stains positive for CK7 and vimentin. CK8 and CK18 in solid tumors. [24] Integrin $\beta 1$, $\beta 2$, $\beta 3$ detected in the pseudocribiform areas. [25] S-100 is almost always present strong and diffuse. [20,26] 80% cells stain positive for WT1. P63 positivity is also noted. 70% cases exhibit PRKD1 mutation. [27]

4.1 Differential diagnosis

PAC is commonly mistaken for Adenoid cystic carcinoma and this is addressed by Ellis & Auclair & Dardick. [8,22] They are mainly differentiated by contour of tumor cells and contour/ nature of luminal spaces and cytology. [21] IHC plays a central role mainly S100 protein, WT1 & SMA like S 100 protein and WT1 are regularly expressed in PAC as opposed to AdCC where SMA is frequently positive.

MYP overexpression is hallmark of AdCC [26] but is absent in PAC [28] High grade transformation has been reported but rare [29].

4.2 Management

Surgical excision provides best locoregional control. [12] SEER database 5-10 year disease specific survival of 98.6 and 96.4% respectively [12] prevention of recurrence is important because recurrent lesion behave far more aggressively. Palatal PAC require extended maxillectomy with rehabilitation. [30] Since metastasis is rare, no need of neck dissection [31]. Radiotherapy and chemotherapy are restricted to palliative cases. [4]

Follow-up of 15-20 years is necessary and chest x-ray for pulmonary metastasis. Prognosis is good in cases with low Ki67 but cases of death are noted.

4. CONCLUSION

Palatal salivary gland malignancies pose a diagnostic dilemma for the pathologists. Although being the second most common salivary gland malignancy, reported cases of PAC are only handful. Sometimes underdiagnosis can also occur due to its bland cytology. Hence any general practitioner should be skeptical about the palatal swelling and refer the patient to a specialist.

REFERENCES

1. Young A, Okuyemi OT. Malignant Tumors of the Palate. In: StatPearls. Treasure Island: StatPearls Publishing. 2022.
2. Batsakis JG, Pinkston GR, Luna MA, et al. Adenocarcinomas of the oral cavity: a clinicopathologic study of terminal duct carcinomas. *J Laryngol Otol*. 1983;97(9):825-35.
3. Freedman PD, Lumerman H. Lobular carcinoma of intraoral minor salivary gland origin: report of twelve cases. *Oral Surg Oral Med Oral Pathol*. 1983;56(2):157-66.

4. Vander Poorten V, Triantafyllou A, Skálová A, Stenman G, Bishop JA, Hauben E, et al. Polymorphous adenocarcinoma of the salivary glands: reappraisal and update. Eur Arch Otorhinolaryngol. 2018;275(7):1681-95.
5. Luna MA, Wenig BM. Polymorphous low-grade adenocarcinoma. In: Barnes L, Eveson JW, Reichart P, Sidransky D, editors. Pathology & Genetics Head and Neck Tumours. Lyon: IARC Press; 2005.
6. Tomar R, Garg N, Agarwal S. Polymorphous low grade adenocarcinoma of lip clinically mimicking squamous cell carcinoma: An unusual presentation. J Cytol. 2015;32(1):59-61.
7. Mills ES, Carter CD. In: Sternberg's Diagnostic Surgical Pathology. Philadelphia: Lippincott Williams & Wilkins; 2010.
8. Ellis G, Auclair P Salivary gland tumors: malignant epithelial tumors. In: Ellis, G, Auclair P Atlas of tumor pathology. Tumors of the salivary glands. Third series, seventeenth fascicle. Armed Forces Institute of Pathology, Washington DC, 1996;155-228.
9. Castle JT, Thompson LDR, Frommelt RA, et al. Poly morphous low grade adenocarcinoma: a clinicopathologic study of 164 cases. Cancer. 1999;86(2):207-19.
10. Skalova A, Sima R, Vanecek T, et al. Acinic cell carcinoma with high-grade transformation: a report of 9 cases with immunohistochemical study and analysis of TP53 and HER-2/neu genes. Am J Surg Pathol. 2009;33(8):1137-45.
11. Seethala RR, Johnson JT, Barnes L, Myers EN. Polymorphous low-grade adenocarcinoma. Arch Otolaryngol Head Neck Surg. 2010;136(4):385-92.
12. Patel TD, Vazquez A, Marchiano E et al. Polymorphous low-grade adenocarcinoma of the head and neck: a populationbased study of 460 cases. Laryngoscope. 2015;125(7):1644-9.
13. Tian Z, Li L, Wang L, et al. Salivary gland neoplasms in oral and maxillofacial regions: a 23-year retrospective study of 6982 cases in an eastern Chinese population. Int J Oral Maxillofac Surg. 2010;39(3):235-242.
14. Bradley PJ, McGurk M. Incidence of salivary gland neoplasms in a defined UK population. Br J Oral Maxillofac Surg. 2013;51(5):399-403.
15. Kato H, Kanematsu M, Makita H, et al. CT and MR imaging findings of palatal tumors. Eur J Radiol. 2014;83(3):e137-46.
16. Hannen EJM, Bulten J, Festen J et al. Polymorphous low grade adenocarcinoma with distant metastases and deletions on chromosome 6q23-qter and 11q23-qter: a case report. J Clin Pathol. 2000;53(12):942-5.
17. Tanaka F, Wada H, Inui K, et al. Pulmonary metastasis of polymorphous low-grade adenocarcinoma of the minor salivary gland. Thorac Cardiovasc Surg 1995;43:178-80.
18. Pfeiffer J, Ridder GJ. Diagnostic value of ultrasoundguided core needle biopsy in patients with salivary gland masses. Int J Oral Maxillofac Surg. 2012;41(4):437-43.
19. Gibbons D, Hossein Saboorian M, Vuitch F, et al. Fine needle aspiration findings in patients with polymorphous low grade adenocarcinoma of the salivary glands. Cancer. 1999;87(1):31-6.

20. Regezi JA, Zarbo RJ, Stewart JCB, Courtney RM. Polymorphous low-grade adenocarcinoma of minor salivary gland. A comparative histologic and immunohistochemical study. Oral Surg Oral Med Oral Pathol. 1991;71(4):469-75.
21. Woolgar J, Triantafyllou A. Contemporary salivary clinical pathology-facts and dilemmas. In: Bradley P, Guntinas-Lichius O Salivary gland disorders and diseases: diagnosis and management, 1st edn. Thieme, Stuttgart, pp 27–41.
22. Dardick I. Color Atlas/text of salivary gland tumor pathology. 1996.
23. Triantafyllou A, Hunt JL, Devaney KO, Ferlito A. A perspective of comparative salivary and breast pathology. Part I: microstructural aspects, adaptations and cellular events. Eur Arch Oto Rhino Laryngol. 2014;271(4):647-63.
24. Marotta Reis de Vasconcellos L, Avila Sarmiento Silveira V, Eduardo Blumer Rosa L et al. Immunohistochemistry as a fundamental tool for the differential diagnosis of polymorphous low-grade adenocarcinoma. Quintessence Int. 2006;37(7):565-73.
25. Araújo V, Sousa S, Jaeger M, et al. Characterization of the cellular component of polymorphous low-grade adenocarcinoma by immunohistochemistry and electron microscopy. Oral Oncol 1999;35(2):164-72.
26. Leader R, Deol-Poonia RK, Sheard J, Triantafyllou A. Immunohistochemical localization of WT1 in epithelial salivary tumors. Pathol Res Pract. 2014;210(11):726-32.
27. Weinreb I, Piscuoglio S, Martelotto LG, et al. Hotspot activating PRKD1 somatic mutations in polymorphous low-grade adenocarcinomas of the salivary glands. Nat Genet. 2014;46(11):1166-9.
28. Persson F, Fehr A, Sundelin K, et al. Studies of genomic imbalances and the MYB-NFIB gene fusion in polymorphous low-grade adenocarcinoma of the head and neck. Int J Oncol. 2012;40(1):80-4.
29. Nagao T. “Dedifferentiation” and high-grade transformation in salivary gland carcinomas. Head Neck Pathol. 2013;7:37-47.
30. El-Naaj IA, Leiser Y, Wolff A, Peled M Polymorphous low grade adenocarcinoma: case series and review of surgical management. J Oral Maxillofac Surg. 2011;69:1967-72.
31. Pogodzinski MS, Sabri AN, Lewis JE, Olsen KD. Retrospective study and review of polymorphous low-grade adenocarcinoma. Laryngoscope. 2006;116(12):2145-9.