

Anogenital Warts in Children Always Means Sexual Abuse?

Mariana Meneses^{1*}, Maria Sousa Dias¹, Sílvia Duarte-Costa¹, Joana Rocha², Maria José Costa^{1,3}

¹Department of Pediatrics, Unidade Local de Saúde de Matosinhos - Hospital Pedro Hispano, Porto, Portugal

²Department of Dermatology, Unidade Local de Saúde de Matosinhos - Hospital Pedro Hispano, Porto, Portugal

³Nucleus to Support Children at risk, Unidade Local de Saúde de Matosinhos - Hospital Pedro Hispano, Porto, Portugal

Citation: Mariana Meneses, Maria Sousa Dias, Sílvia Duarte-Costa, Joana Rocha, Maria José Costa. Anogenital Warts in Children Always Means Sexual Abuse?. *Int Clin Med Case Rep Jour.* 2024;3(11):1-3.

Received Date: 07 November, 2024; **Accepted Date:** 08 November, 2024; **Published Date:** 10 November, 2024

***Corresponding author:** Mariana Meneses, ¹Department of Pediatrics, Unidade Local de Saúde de Matosinhos - Hospital Pedro Hispano, Porto, Portugal

Copyright: ©2024 Mariana Meneses. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

CLINICAL IMAGE



Figure 1 Condyloma acuminatum before treatment




Figure 2 Condyloma acuminatum after treatment

Previously healthy 8-year-old female patient with multiple anogenital lesions with “cauliflower appearance”, which were increasing in size, and causing some pain (**Figure 1**). After assessment, the diagnosis of condyloma acuminatum was made and treatment with imiquimod was started. The child lived with her mother and older sister, and neither of them had condylomatosis. Facing the positivity of HPV infection, analytical study to exclude other sexually transmitted diseases was made (negative). Psychologic and forensic evaluation concluded that the child was stable at a behavioral and emotional level, with no signs of sexual abuse. Due to persistence of lesions under imiquimod, cryotherapy was started with a subsequent improvement (**Figure 2**).

Anogenital warts are manifestations of HPV infection.^[1] Epidemiologic data in children are limited and the exact prevalence of this condition is unknown. A female predominance is suggested by several studies.^[2,3,4] In children, the serotypes detected are variable and the possibility of sexual abuse is a major concern^{1,2}. Nevertheless, other modes of viral transmission may account for most of pediatric cases.^[1,2,6] The identification of the virus transmission route is hindered by the HPV variable incubation period.^[6] The diagnosis is made via clinical examination.¹ Viral serotyping isn't routinely performed since it's not necessary for diagnosis and is not reliable for identifying the source of infection.^[2,3,4,8] Interviews with the caregivers and the child, as well as clinical and laboratory testing to evaluate for sexual abuse and other sexually transmitted infections are important parts of the patient assessment.^[1,8] If any signs of sexual abuse are noted, the case must be reported to child protective services according with local policies.^[7]

With this case the authors pretend to enhance that the origin of pediatric anogenital HPV infections remains often untraced, with no indication of sexual abuse, although this possibility should always be considered.

REFERENCES

1. Mammas IN, Sourvinos G, Spandidos DA. Human papilloma virus (HPV) infection in children and adolescents. Eur J Pediatr. 2009;168(3):267-273.
2. Marcoux D et al. Pediatric anogenital warts: a 7-year review of children referred to a tertiary-care hospital in Montreal, Canada. Pediatr Dermatol. 2006;23:199-207.
3. Handley JM et al. Anogenital warts in children. Clin Exp Dermatol 1993;18:241-247.

4. Obalek S et al. Anogenital warts in children. Clin Dermatol. 1997; 15:369-376.
5. Cao CD, Merjanian L, Pierre J, Balica A. A Discussion of High-Risk HPV in a 6-Year-Old Female Survivor of Child Sexual Abuse. Case Rep Obstet Gynecol. 2017;2017:1-4.
6. Syrjänen S. Current concepts on human papillomavirus infections in children. Apmis. 2010;118(6-7):494-509.
7. Rock B, Naghashfar Z, Barnett N, Buscema J, Woodruff JD, Shah K. Genital Tract Papillomavirus Infection in Children. Arch Dermatol. 1986;122(10):1129-1132.
8. Unger ER, Fajman NN, Maloney EM, et al. Anogenital human papillomavirus in sexually abused and nonabused children: A multicenter study. Pediatrics. 2011;128(3).