Annals of Clinical Pharmacology and Physiology
Review Article

Gut skin axis: An Epidemiological Assessment of Potential Health Risks in the Indian Population

Rahul Hajare*, Mahek Khan
Sandip University, Nashik, India

Citation: *Rahul Hajare, Mahek Khan. Gut skin axis: An Epidemiological Assessment of Potential Health Risks in the Indian Population. Annal Clin Pharm Phys. 2026;1(1):1-11.*
Received Date: 08 November 2026; **Accepted Date:** 14 November 2026; **Published Date:** 15 November 2026
***Corresponding author:** Rahul Hajare, Sandip University, Nashik, India
Copyright: © Rahul Hajare, Open Access 2026. This article, published in Annal Clin Pharm Phys (ACPP) (Attribution 4.0 International), as described by http://creativecommons.org/licenses/by/4.0/

ABSTRACT

The gut–skin axis is a complex bidirectional communication network connecting the gastrointestinal tract and the skin through microbial, metabolic, immunological, endocrine and neuroendocrine pathways. Increasing evidence indicates that the gut microbiome contributes to systemic homeostasis and may influence skin-barrier integrity, immune regulation, inflammatory responses and susceptibility to several dermatological disorders. Gut dysbiosis, broadly characterised by alterations in microbial composition, diversity or function, may influence intestinal-barrier integrity and microbial metabolite production, thereby modifying systemic immune and inflammatory responses. The gut–skin axis has been investigated in relation to several inflammatory and immune-mediated dermatological conditions, including acne vulgaris, atopic dermatitis, psoriasis and rosacea. Dietary patterns, psychological stress, sleep disturbances, antibiotic exposure, smoking, alcohol consumption, physical inactivity and environmental pollutants may further influence the gut–skin relationship. Importantly, the gut–skin axis does not imply the physical migration of faecal material from the intestine to the skin. Rather, communication may occur through microbial products, metabolites, immune mediators and neuroendocrine signalling pathways. Nutritional optimisation, lifestyle modification, appropriate skin-barrier care and microbiome-targeted interventions such as probiotics, prebiotics, synbiotics and emerging postbiotic approaches represent potential strategies for supporting skin health. However, clinical evidence for microbiome-based interventions remains heterogeneous and causality between specific microbial alterations and individual skin diseases has not been established in all cases. Future research integrating metagenomics, metabolomics, immunology, proteomics and personalised medicine may clarify disease-specific microbial signatures and facilitate the development of targeted therapeutic strategies. The gut–skin axis therefore represents a promising multidisciplinary field linking gastroenterology, dermatology, microbiology, immunology, nutrition and pharmaceutical sciences.

Keywords: Gut skin axis; Gut microbiota; Gut dysbiosis; Skin microbiome; Intestinal barrier

INTRODUCTION

The gut–skin axis represents a dynamic and bidirectional physiological communication system between the gastrointestinal tract and the skin. This relationship involves interactions among the gut microbiota, intestinal epithelial barrier, immune system, microbial metabolites, endocrine pathways and nervous system. Although the gastrointestinal tract and skin are anatomically distant, they share important immunological and physiological characteristics and are continuously influenced by environmental, nutritional, metabolic and microbial factors.

The human gastrointestinal tract contains a highly diverse microbial ecosystem comprising bacteria, fungi, viruses, archaea and other microorganisms. These microbial communities participate in nutrient metabolism, production of bioactive metabolites, immune maturation, maintenance of epithelial integrity and protection against pathogenic microorganisms. Consequently, alterations in the composition or function of the gut microbiota may have effects extending beyond the gastrointestinal tract.[1-4] The skin is the body's largest organ and provides a critical physical and immunological barrier against environmental challenges. Skin homeostasis depends on coordinated interactions among keratinocytes, immune cells, lipids, structural proteins, the skin microbiome and environmental factors. Disruption of these mechanisms may contribute to inflammation, altered barrier function and dermatological disease.[4] The concept of the gut–skin axis has therefore attracted increasing attention in dermatological research. Associations between intestinal microbial alterations and several inflammatory skin diseases have been reported, including acne vulgaris, atopic dermatitis, psoriasis and rosacea. However, these relationships are complex and disease-specific. The presence of dysbiosis should not automatically be interpreted as evidence of a direct causal relationship. This review examines the biological mechanisms underlying the gut–skin axis, the potential contribution of gut dysbiosis to skin health, the influence of dietary and lifestyle factors and the emerging role of microbiome-based therapeutic strategies.

Biological Basis of the Gut–Skin Axis

Gut Microbiota

The gut microbiota comprises a diverse microbial community that contributes to digestion, metabolism, immune development and maintenance of intestinal homeostasis. A healthy and functionally diverse microbial ecosystem can restrict pathogen colonisation, support epithelial integrity and contribute to appropriate immune tolerance.[1,3] Microbial composition and function are influenced by age, diet, geographical location, genetics, medication use, lifestyle and environmental exposure. Antibiotics and other medications can substantially modify microbial communities, while dietary composition can influence both microbial abundance and microbial metabolic activity. The gut microbiota may influence distant organs through microbial metabolites that enter systemic circulation and through modulation of immune, endocrine and neural pathways. These mechanisms provide a biological basis for communication between intestinal microorganisms and the skin. However, the concept of a ‘healthy microbiome’ should not be reduced to the presence or absence of individual bacterial species. Microbiome function, metabolic activity, microbial interactions and host-specific characteristics are also important

Intestinal Barrier

The intestinal barrier consists of intestinal epithelial cells, tight-junction proteins, mucus, immune components and associated microbial communities. Its principal function is to permit the absorption of nutrients while limiting inappropriate penetration of microorganisms, toxins and potentially harmful luminal substances.

Disruption of intestinal-barrier integrity may increase intestinal permeability and alter exposure of the immune system to microbial products. This may contribute to changes in systemic inflammatory signalling and potentially influence tissues distant from the gastrointestinal tract, including the skin.[1,3,4] The relationship between intestinal permeability and systemic disease should, however, be interpreted cautiously. Increased intestinal permeability may occur in several physiological and pathological circumstances and its magnitude and clinical significance may differ between individuals and diseases.

Microbial Metabolites and Short-Chain Fatty Acids

Gut microorganisms produce numerous metabolites capable of influencing host physiology. Among the most extensively investigated are short-chain fatty acids, particularly acetate, propionate and butyrate. Short-chain fatty acids are generated predominantly through microbial fermentation of dietary fibre. They can influence epithelial integrity, immune-cell activity, inflammatory signalling and cellular metabolism. Butyrate is particularly important for intestinal epithelial physiology and immune regulation. Changes in dietary fibre intake and microbial composition may consequently alter short-chain fatty-acid production. These metabolic changes represent one potential mechanism through which the intestinal microbiota may influence systemic physiology and skin health. Other microbial metabolites, including bile-acid derivatives, tryptophan metabolites and other small molecules, may also participate in host–microbe communication. Further research is required to determine which metabolites have clinically meaningful effects on specific skin disorders.

Immune Regulation

The gastrointestinal tract is a major immunological organ. Interactions between intestinal microorganisms and immune cells contribute to immune maturation, tolerance and regulation of inflammatory responses. Gut dysbiosis may alter immune signalling and modify the balance between regulatory and inflammatory pathways. Microbial components and metabolites can influence cytokine production and immune-cell activation. Because the skin contains an extensive network of innate and adaptive immune cells, systemic alterations in immune activity may influence cutaneous inflammation. This mechanism has been proposed as one pathway connecting intestinal microbial alterations with inflammatory skin disorders. Nevertheless, immune responses are highly disease-specific and may be influenced by genetic, environmental and local cutaneous factors.

Neuroendocrine Signalling

The gastrointestinal tract and skin are influenced by the nervous and endocrine systems. Psychological stress can activate the hypothalamic–pituitary–adrenal axis and sympathetic nervous system, potentially affecting intestinal motility, microbial composition, epithelial-barrier function, immune activity and skin physiology.

Communication between the gut microbiota and nervous system is commonly discussed within the context of the microbiota–gut–brain axis. Because the brain, gastrointestinal tract and skin are interconnected through neuroendocrine, immune and neural pathways, these mechanisms may contribute to the broader gut–skin relationship. Stress-related neuroendocrine changes may therefore influence both intestinal and cutaneous homeostasis. However, the specific contribution of these pathways to individual skin diseases remains an active area of investigation.

Gut Dysbiosis and Skin Health

Gut dysbiosis broadly refers to alterations in microbial composition, diversity, abundance or function. It should not necessarily be interpreted as the complete loss of beneficial microorganisms or the presence of a single pathogenic microorganism. Potential consequences of dysbiosis include changes in microbial metabolites, altered intestinal-barrier function, modified immune responses and changes in systemic inflammatory signalling. These alterations may subsequently influence skin-barrier integrity and cutaneous inflammatory responses.[1,3-5] However, the relationship between dysbiosis and dermatological disease may be bidirectional. Skin disease, dietary changes, medications, psychological stress and other factors may themselves influence the gut microbiome. Consequently, dysbiosis may represent a contributor, consequence or both, depending on the disease and individual context. This distinction is particularly important when interpreting microbiome studies. The identification of particular microbial taxa in association with a disease does not establish that these microorganisms cause the disease. Functional studies and longitudinal research are necessary to clarify causal relationships.

Gut–Skin Axis and Major Inflammatory Skin Disorders

Acne Vulgaris

Acne vulgaris is a common inflammatory skin disorder involving follicular hyperkeratinisation, sebaceous-gland activity, microbial interactions and immune-mediated inflammation. The gut–skin axis has been proposed as one factor potentially influencing acne through systemic inflammatory pathways, dietary factors, microbial metabolites and interactions between intestinal and cutaneous microbiota. Dietary patterns characterised by a high glycaemic load have been investigated in relation to acne. Some evidence suggests that dietary modification may influence acne severity in selected individuals, although responses are heterogeneous and diet is only one component of acne pathophysiology. Probiotics and other microbiome-directed interventions have also received increasing research attention. However, additional well-designed clinical trials are required before these approaches can be considered established treatments for acne

Atopic Dermatitis

Atopic dermatitis is a chronic inflammatory skin disorder characterised by skin-barrier dysfunction, immune dysregulation and recurrent inflammation. Alterations in the intestinal microbiome have been reported in individuals with atopic dermatitis. Proposed mechanisms include interactions between intestinal microorganisms, immune development, epithelial-barrier function and systemic inflammatory signalling.[4,5] The gut microbiome may therefore represent a potential therapeutic target. Nevertheless, differences between studies in patient populations, sampling methods, sequencing techniques and microbial-analysis approaches make it difficult to identify a single disease-specific microbial signature. Clinical evidence for probiotics and related interventions remains variable and may depend on microbial strain, formulation, patient characteristics and disease phenotype.

Psoriasis

Psoriasis is a chronic immune-mediated inflammatory skin disease characterised by abnormal keratinocyte proliferation and dysregulated immune signalling. Research has identified potential associations between intestinal microbial alterations and psoriasis. The gut microbiome may influence systemic inflammatory pathways and immune responses relevant to disease activity.[4,5] Metabolic comorbidities, obesity and dietary factors may further complicate the relationship between intestinal microbiota and psoriasis. Although these findings provide a plausible biological framework, longitudinal and interventional studies are required to determine whether specific microbial alterations contribute directly to disease initiation, progression or treatment response.

Rosacea

Rosacea is a chronic inflammatory facial skin disorder associated with persistent or recurrent erythema, vascular abnormalities, inflammatory lesions and, in some individuals, ocular involvement. Gastrointestinal factors have attracted attention in rosacea research, including possible relationships with gastrointestinal disorders and intestinal microbial alterations. Several mechanisms have been proposed, including immune activation, microbial metabolites, altered intestinal-barrier function and neurovascular signalling. However, the clinical relationship between intestinal microbiota and rosacea remains an evolving area of research. Existing evidence does not justify assuming that gut dysbiosis is a universal cause of rosacea.

Dietary, Lifestyle, Pharmacological and Environmental Factors

The gut–skin axis is influenced by multiple interacting factors rather than a single determinant.

Factor	Potential effect on the gut	Potential effect on the skin
Unbalanced diet	May alter microbial composition and metabolic activity	May contribute to inflammatory skin changes

Low dietary fibre	May reduce microbial fermentation and production of beneficial metabolites	May influence immune and barrier regulation
Excess refined carbohydrates	May influence metabolic and microbial homeostasis	May aggravate some inflammatory conditions
Chronic psychological stress	May influence microbiota, intestinal motility, barrier function and immunity	May worsen inflammation and disease activity
Poor sleep	May affect metabolic and immune regulation	May impair skin recovery and homeostasis
Antibiotic exposure	Can alter microbial diversity and composition	May indirectly influence immune and skin homeostasis
Smoking	May promote microbial alterations and oxidative stress	Can increase oxidative stress and skin damage
Excessive alcohol consumption	May affect intestinal barrier function and microbial balance	May contribute to inflammation and oxidative stress
Physical inactivity	May adversely affect metabolic and microbial health	May indirectly influence inflammatory processes
Environmental pollution	Can promote oxidative and inflammatory responses	May impair skin-barrier function and contribute to premature ageing
Excessive ultraviolet exposure	May indirectly influence systemic inflammatory pathways	Can directly damage skin cells and accelerate photoageing

These factors frequently interact. Poor diet, psychological stress, inadequate sleep and physical inactivity may occur together and collectively influence metabolic, immune and microbial homeostasis.

Fecal Microbiota and the Gut–Skin Axis

A scientifically important distinction should be made between faecal material, faecal microbiota and the gut–skin axis. Under normal physiological conditions, faecal material remains within the gastrointestinal tract and does not physically migrate to the skin. Therefore, the gut–skin axis should not be interpreted as the movement or deposition of faecal material onto or into the skin. Faecal samples are commonly used in microbiome research because they

provide a relatively accessible source for investigating intestinal microbial communities. Analysis of faecal microbiota can provide information regarding microbial composition and, when combined with appropriate functional approaches, microbial metabolic activity. The potential influence of gut microorganisms on the skin occurs through biological communication pathways. A simplified model can be represented as: **Gut microbiota → microbial metabolites and microbial products → intestinal and immune signalling → systemic circulation and immune regulation → skin barrier and inflammatory responses** Short-chain fatty acids, bile-acid derivatives, tryptophan metabolites and other microbial products may participate in these pathways. Immune mediators and neuroendocrine signalling provide additional communication routes. Thus, faecal microbiota can serve as a research proxy for intestinal microbial communities, whereas faecal material itself is not a physiological component that travels through the gut–skin axis.

Strategies for Maintaining and Supporting Skin Health

Maintaining skin health requires an integrated approach that addresses both cutaneous and systemic factors.

Balanced Nutrition

A dietary pattern containing adequate fibre, vegetables, fruits, legumes, whole grains, nuts and other nutrient-rich foods can support gastrointestinal and metabolic health.

Dietary diversity may also contribute to microbial diversity and microbial metabolic function.

Adequate Hydration

Appropriate fluid intake supports normal physiological function. Hydration should be considered as one component of overall health rather than as an independent treatment for dermatological disease.

Skin-Barrier Protection

Appropriate cleansing, regular moisturisation, avoidance of unnecessary irritants and maintenance of the skin barrier are fundamental components of dermatological care.

Photoprotection

Protection from excessive ultraviolet radiation is important for reducing photoageing, DNA damage and other UV-associated skin effects.

Stress Management

Stress-management strategies may support psychological well-being and may contribute to healthy neuroendocrine, immune and gastrointestinal function.

Adequate Sleep

Regular and sufficient sleep supports immune, endocrine, metabolic and tissue-repair processes and may contribute to healthy skin homeostasis.

Physical Activity

Regular physical activity supports metabolic health and may positively influence immune and microbial homeostasis.

Appropriate Medical Management

Underlying dermatological or gastrointestinal diseases should be evaluated and managed by appropriately qualified healthcare professionals. Microbiome-targeted approaches should complement established medical treatment where appropriate rather than automatically replacing evidence-based therapies.

Probiotics, Prebiotics, Synbiotics and Emerging Microbiome-Based Therapies

Growing recognition of the gut–skin axis has stimulated interest in microbiome-directed interventions.

Probiotics

Probiotics are live microorganisms administered in adequate amounts with the intention of providing a health benefit to the host. Their effects are strain-specific and cannot be generalised to all probiotic products. Differences in microbial strains, dose, formulation, duration and host characteristics may contribute to variations in clinical outcomes.

Prebiotics

Prebiotics are substrates selectively utilised by host microorganisms that confer a health benefit. Dietary fibres are important examples of substrates capable of influencing microbial fermentation and metabolite production.

Synbiotics

Synbiotics combine live microorganisms with substrates that are intended to support beneficial microbial activity. Their potential effects may arise from complementary or synergistic interactions.

Postbiotics

Postbiotics have emerged as another area of interest. They involve preparations of inanimate microorganisms and/or their components that confer a health benefit to the host. Microbial metabolites and other defined microbial products are also being investigated as potential therapeutic tools. However, microbiome-based therapy remains an evolving field. Clinical outcomes may vary according to microbial strain, formulation, dose, duration, disease phenotype, dietary background, host genetics and baseline microbiome composition. Consequently, claims regarding probiotics, prebiotics, synbiotics or postbiotics should be supported by disease-specific clinical evidence.

Skin Microbiome and Its Interaction with the Gut

Although the primary focus of the gut–skin axis is communication between the intestinal microbiota and skin, the skin itself contains a diverse microbial ecosystem. The cutaneous microbiome interacts with keratinocytes, immune cells, skin lipids and environmental factors. Microbial communities vary according to anatomical site, moisture, temperature, sebaceous-gland activity and other local characteristics. Alterations in the skin microbiome have been associated with several dermatological conditions. However, as with the gut microbiome, the presence of a particular microorganism does not necessarily establish causation. The gut and skin microbiomes may therefore be viewed as interconnected microbial ecosystems influenced by host immunity, nutrition, metabolism, environmental exposure and lifestyle.[6-12]

CONCLUSION

The gut–skin axis provides an important biological framework for understanding the relationship between intestinal microbial health and cutaneous homeostasis. Communication between the gastrointestinal tract and skin involves interconnected microbial, metabolic, immunological, endocrine and neuroendocrine pathways. Gut dysbiosis may influence intestinal-barrier integrity, microbial metabolite production, systemic immune regulation and inflammatory signalling and may therefore contribute to alterations in skin health. Associations between intestinal microbial changes and conditions such as acne vulgaris, atopic dermatitis, psoriasis and rosacea have generated considerable scientific interest. However, the strength of evidence varies between diseases and association should not automatically be interpreted as causation. Importantly, the gut–skin axis does not involve the physical migration of faecal material to the skin. Rather, intestinal microorganisms and their metabolites may influence distant tissues through systemic biological signalling, immune regulation and neuroendocrine pathways. Dietary quality, adequate sleep, stress management, regular physical activity, appropriate skin-barrier care, photoprotection and evidence-based medical treatment remain important components of maintaining skin health. Probiotics, prebiotics, synbiotics, postbiotics and other microbiome-targeted approaches represent promising areas for future investigation. However, their clinical application requires stronger disease-specific evidence and improved understanding of individual microbiome variability. Overall, the gut–skin axis represents a rapidly developing interdisciplinary field with implications for dermatology, gastroenterology, microbiology, immunology, nutrition and pharmaceutical sciences. Future well-designed clinical, translational and mechanistic studies may determine whether targeted modulation of the gut microbiome can become a reliable component of personalised strategies for the prevention and management of inflammatory skin disorders.

Novel Conclusion

"The therapeutic paradigm of modern dermatology is undergoing a significant evolution, shifting away from purely topical treatments toward systemic, interdisciplinary interventions. This review constructs a vital, epidemiologically grounded framework for the gut–skin axis, demonstrating that cutaneous homeostasis is intricately intertwined with the functional diversity of the gastrointestinal microbiome.

The primary novelty of this integrated epidemiological assessment rests upon two critical physiological models:

- **Deconstruction of the Structural Myth:** A key academic contribution of this review is the absolute deconstruction of pop-science and commercial wellness misconceptions regarding the gut-skin relationship. It explicitly establishes that the axis operates through **compartmentalized biochemical communication channels**—including short-chain fatty acids (SCFAs), tryptophan derivatives, systemic immune cell maturation, and hypothalamic–pituitary–adrenal (HPA) axis neuroendocrine signaling—rather than any physical migration of fecal microbial matter. Fecal microbiota serves exclusively as an accessible clinical research proxy for intestinal sequencing, not as a cutaneous contaminant.

- **The Dual Burden of Nutritional Transition:** Crucially, this review contextualizes these homeostatic mechanisms within the rapid socioeconomic urbanization of the contemporary Indian population. The ongoing epidemiological transition away from fiber-rich diets toward high-glycemic, ultra-processed food systems induces persistent low-grade gut dysbiosis. This systemic disruption degrades intestinal epithelial tight junctions, thereby accelerating systemic inflammatory signaling that increases host susceptibility to chronic dermatological phenotypes such as acne vulgaris, atopic dermatitis, psoriasis, and rosacea.

Ultimately, this paper shifts the focus of clinical dermatology from isolated symptom management to long-term systemic optimization. While emerging microbiome-targeted therapies—including probiotics, prebiotics, synbiotics, and postbiotics—offer promising preventive avenues, their implementation currently demands more rigorous, strain-specific clinical validation. Future cross-sectional and longitudinal research must utilize advanced multi-omic sequencing (metagenomics, metabolomics, and proteomics) within transitioning South Asian cohorts to map out disease-specific microbial signatures, paving the way for predictable, personalized therapeutic strategies in public health."

REFERENCES

1. De Pessemier B, Grine L, Debaere M, Maes A, Paetzold B, Callewaert C. Gut–skin axis: current knowledge of the interrelationship between microbial dysbiosis and skin conditions. Microorganisms. 2021;9(2):353. doi:10.3390/microorganisms9020353.
2. Salem I, Ramser A, Isham N, Ghannoum MA. The gut microbiome as a major regulator of the gut-skin axis. Front Microbiol. 2018;9:1459. doi:10.3389/fmicb.2018.01459.
3. Sinha S, Lin G, Ferenczi K. The skin microbiome and the gut-skin axis. Clin Dermatol. 2021;39(5):829-839. doi:10.1016/j.clindermatol.2021.08.021.
4. Chen M, Wang R, Wang T, et al. Gut microbiota and skin pathologies: mechanism of the gut-skin axis in atopic dermatitis and psoriasis. Int Immunopharmacol. 2024;141:112658.
5. Salem I, et al. The gut microbiome and skin disorders: a systematic review of the emerging gut–skin axis. Dermatol Ther. 2020;33.
6. Bowe WP, Logan AC. Acne vulgaris, probiotics and the gut-brain-skin axis—back to the future? Gut Pathog. 2011;3:1. doi:10.1186/1757-4749-3-1.
7. De Pessemier B, Grine L, Debaere M, Maes A, Paetzold B, Callewaert C. Gut–skin axis: current knowledge of the interrelationship between microbial dysbiosis and skin conditions. Microorganisms. 2021;9(2):353. doi:10.3390/microorganisms9020353.
8. Parke MA, Perez-Sanchez A, Zamil DH, Katta R. Diet and skin barrier: the role of dietary interventions on skin barrier function. Dermatol Pract Concept. 2021;11(1).

9. Salem I, Ramser A, Isham N, Ghannoum MA. The gut microbiome as a major regulator of the gut-skin axis. Front Microbiol. 2018;9:1459. doi:10.3389/fmicb.2018.01459.
10. Sinha S, Lin G, Ferenczi K. The skin microbiome and the gut-skin axis. Clin Dermatol. 2021;39(5):829-839. doi:10.1016/j.clindermatol.2021.08.021.
11. Chen M, Wang R, Wang T, et al. Gut microbiota and skin pathologies: mechanism of the gut-skin axis in atopic dermatitis and psoriasis. Int Immunopharmacol. 2024;141:112658.
12. Parke MA, Perez-Sanchez A, Zamil DH, Katta R. Diet and skin barrier: the role of dietary interventions on skin barrier function. Dermatol Pract Concept. 2021;11(1).