




**VOQUEZNA Triple Pak**<sup>®</sup>  
vonoprazan tablets 20 mg  
amoxicillin capsules 500 mg  
clarithromycin tablets 500 mg



**VOQUEZNA Dual Pak**<sup>®</sup>  
vonoprazan tablets 20 mg  
amoxicillin capsules 500 mg

# THE POWER TO HELP FIGHT *H. PYLORI* INFECTION

Learn about *H. pylori* and  
a unique 14-day treatment



Find out  
about savings  
options  
(pages 10 and 11)

## INDICATION AND IMPORTANT SAFETY INFORMATION

### What are VOQUEZNA TRIPLE PAK and VOQUEZNA DUAL PAK?


- VOQUEZNA<sup>®</sup> TRIPLE PAK<sup>®</sup> (vonoprazan tablets, amoxicillin capsules, clarithromycin tablets) and VOQUEZNA<sup>®</sup> DUAL PAK<sup>®</sup> (vonoprazan tablets, amoxicillin capsules) are co-packaged prescription medicines for the treatment of a *Helicobacter pylori* (*H. pylori*) bacterial infection in adults.
- It is not known if VOQUEZNA TRIPLE PAK and VOQUEZNA DUAL PAK are safe and effective in children.
- To reduce the development of drug-resistant bacteria and maintain the effectiveness of VOQUEZNA TRIPLE PAK, VOQUEZNA DUAL PAK and other antibacterial drugs, these products should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria.

### Do not take VOQUEZNA TRIPLE PAK if you:

- are known to have an allergy or be sensitive to the components of VOQUEZNA TRIPLE PAK (vonoprazan, amoxicillin, clarithromycin), macrolide antibiotics (such as azithromycin and erythromycin), or penicillin.

Please see additional Important Safety Information throughout and full Prescribing Information.

## ESSENTIAL INFORMATION ABOUT *H. PYLORI* INFECTION AND TREATMENT



### *H. pylori* is a bacterial infection that occurs in the stomach

The infection may cause symptoms like pain or a burning sensation in the stomach, nausea, bloating, loss of appetite, and frequent burping.



*H. pylori* should be treated when detected whether there are symptoms or not, as it is contagious. Leaving *H. pylori* untreated increases the risk of related health issues.

### *H. pylori* is common

It's estimated that 50% of the world's population and 36% of the US population have *H. pylori* infection.



### IMPORTANT SAFETY INFORMATION (cont'd)

**Do not take VOQUEZNA TRIPLE PAK if you:**

- are taking:
  - medicines that contain rilpivirine (EDURANT, COMPLERA, JULUCA, ODEFSEY)
  - pimozide
  - colchicine, if you have kidney or liver problems
  - lomitapide, lovastatin, and simvastatin
  - ergot alkaloids (ergotamine or dihydroergotamine)
  - lurasidone
- have a history of yellowing of the skin (jaundice) or liver problems when taking clarithromycin.
- are pregnant or plan to become pregnant except in clinical circumstances where your healthcare provider has determined no alternative therapy is appropriate. Clarithromycin, a medicine in VOQUEZNA TRIPLE PAK may harm your unborn baby.

### Leaving *H. pylori* untreated can lead to health problems

These health concerns include stomach irritation and swelling, sores in the protective lining of the stomach called ulcers, and an increased risk of certain types of stomach cancers.



### It's important to complete your VOQUEZNA-based 14-day treatment

Taking all of the medication prescribed by your doctor as directed helps increase the chance of getting rid of your *H. pylori* infection.



If you experience side effects that concern you, ask your doctor if you should stop treatment.

Learn more about what makes your treatment unique at [VOQUEZNA.com/HP](https://www.voquezna.com/HP)


### IMPORTANT SAFETY INFORMATION (cont'd)

**Do not take VOQUEZNA DUAL PAK if you:**

- are known to have an allergy or be sensitive to the components of VOQUEZNA DUAL PAK (vonoprazan, amoxicillin) or penicillin.
- are taking medicines that contain rilpivirine (EDURANT, COMPLERA, JULUCA, ODEFSEY).

Please see additional Important Safety Information throughout and full [Prescribing Information](#).

## YOUR *H. PYLORI* TREATMENT REGIMEN CONTAINS ANTIBIOTICS AND AN ACID BLOCKER, VONOPRAZAN



### An effective acid blocker is important for successful treatment

Acid blockers make the stomach less acidic which helps *H. pylori* treatments work better. In your VOQUEZNA-based regimen, vonoprazan is the acid blocker.



### Vonoprazan works quickly to reduce stomach acid

In a study of healthy adults, vonoprazan began blocking acid within 2 to 3 hours.



### Vonoprazan can be taken with or without food

Unlike some other acid blockers, there are no food requirements with vonoprazan.



### IMPORTANT SAFETY INFORMATION (cont'd)

**Before you take VOQUEZNA TRIPLE PAK or VOQUEZNA DUAL PAK, tell your healthcare provider about all of your medical conditions, including if you:**

- are pregnant, think you may be pregnant, or plan to become pregnant. It is not known if VOQUEZNA will harm your unborn baby.
  - **Pregnancy Exposure Registry:** There is a pregnancy registry for women who take VOQUEZNA TRIPLE PAK or VOQUEZNA DUAL PAK during pregnancy. The purpose of this registry is to collect information about the health of you and your baby. If you are pregnant or become pregnant during treatment with VOQUEZNA TRIPLE PAK or VOQUEZNA DUAL PAK, talk to your healthcare provider about how you can join this pregnancy registry or you can enroll in the registry by calling 1-866-609-1612 or visiting <https://voqueznapregnancyregistry.com/>.

### Acid blockers help antibiotics work better

Antibiotics kill *H. pylori* bacteria to clear the infection. They work better when there is less acid in the stomach. Taking an acid blocker, like vonoprazan, along with antibiotics, improves the chance of clearing the infection.



These VOQUEZNA-based regimens are FDA-approved medications

### IMPORTANT SAFETY INFORMATION (cont'd)

**Before you take VOQUEZNA TRIPLE PAK or VOQUEZNA DUAL PAK, tell your healthcare provider about all of your medical conditions, including if you:**

- are breastfeeding or plan to breastfeed. You and your healthcare provider should decide if you will take VOQUEZNA TRIPLE PAK or VOQUEZNA DUAL PAK while breastfeeding.
- have kidney or liver problems.
- have myasthenia gravis.

**Tell your healthcare provider about all the medications you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements. There can be serious side effects when some are used in combination with this product. Serious adverse reactions can occur with VOQUEZNA TRIPLE PAK due to drug interactions of clarithromycin with colchicine, some lipid lowering agents, some calcium channel blockers, and other drugs.

**Please see additional Important Safety Information throughout and full [Prescribing Information](#).**

## FDA-APPROVED TREATMENT REGIMENS TO HELP YOU CLEAR THE INFECTION



### Proven in a clinical trial to get rid of *H. pylori* infection

In a clinical trial, VOQUEZNA® Triple Pak® and VOQUEZNA® Dual Pak® were proven to get rid of *H. pylori* infection for more than 77% of patients after one 14-day treatment (compared to 69% on lansoprazole, amoxicillin, and clarithromycin).



### Possible side effects of your prescription

In a clinical trial, common side effects included: diarrhea, altered sense of taste, vaginal yeast infection, stomach pain, headache, high blood pressure, and cold-like symptoms.

Other more serious side effects are possible. See Important Safety Information included throughout this brochure.

### IMPORTANT SAFETY INFORMATION (cont'd)

#### What are the possible side effects of VOQUEZNA TRIPLE PAK and VOQUEZNA DUAL PAK?

**VOQUEZNA TRIPLE PAK or VOQUEZNA DUAL PAK can cause serious side effects including:**

- **Allergic reactions (hypersensitivity):** Call your doctor right away if you have rash, hives, or other skin changes, face swelling or difficulty breathing.
- **A type of kidney problem (acute tubulointerstitial nephritis):** Some people who take vonoprazan, a component of VOQUEZNA TRIPLE PAK or VOQUEZNA DUAL PAK, may develop a kidney problem called acute tubulointerstitial nephritis. Call your healthcare provider right away if you have a decrease in the amount that you urinate or if you notice blood in your urine.
- **Severe skin reactions:** VOQUEZNA TRIPLE PAK and VOQUEZNA DUAL PAK can cause severe skin reactions, such as skin rash or allergic reaction on or in any part of your body. Symptoms can also include, but are not limited to, fever, chills, body aches or shortness of breath. If you experience any of these symptoms, stop taking VOQUEZNA TRIPLE PAK and VOQUEZNA DUAL PAK immediately and call your doctor right away.



“About a month after completing my 14-day treatment, my doctor tested me and confirmed that the infection was gone. I hope it can work for you as well as it worked for me!”

*Individual results may vary.*

— LaDawn,  
Clinical Trial Patient for VOQUEZNA® Dual Pak®

### IMPORTANT SAFETY INFORMATION (cont'd)

#### What are the possible side effects of VOQUEZNA TRIPLE PAK and VOQUEZNA DUAL PAK?

**VOQUEZNA TRIPLE PAK or VOQUEZNA DUAL PAK can cause serious side effects including:**


- **Gastrointestinal hypersensitivity (drug-induced enterocolitis syndrome):** A type of allergic reaction that causes inflammation in the small and large intestine has been reported with use of amoxicillin, a component of VOQUEZNA TRIPLE PAK and VOQUEZNA DUAL PAK, mostly in patients ≤18 years of age. Symptoms include vomiting occurring 1 to 4 hours after taking the drug and may also be associated with skin paleness, tiredness, low blood pressure, shock, and diarrhea within 24 hours after taking amoxicillin. If you experience any of these symptoms, stop taking VOQUEZNA TRIPLE PAK or VOQUEZNA DUAL PAK and notify your doctor.
- **Severe diarrhea:** Call your doctor right away if you have watery stool, stomach pain, and fever that does not go away while taking VOQUEZNA TRIPLE PAK or VOQUEZNA DUAL PAK or after therapy is completed.

Please see additional Important Safety Information throughout and full [Prescribing Information](#).

# YOUR DOCTOR PRESCRIBED ONE OF THESE VOQUEZNA-BASED OPTIONS FOR YOU



Take your medication 2 times per day: in the morning and in the evening, 12 hours apart.



Pills not actual size.

**14-day treatment** that can be taken with or without food

### IF YOU MISS A DOSE OF VOQUEZNA® TRIPLE PAK®

Take the next dose as soon as possible unless 4 hours have passed since you were supposed to take it. If more than 4 hours have passed, skip the missed dose and take the next dose on time. Continue to take your medicine until all doses in the pack have been taken, even if symptoms go away.

## IMPORTANT SAFETY INFORMATION (cont'd)


What are the possible side effects of VOQUEZNA TRIPLE PAK and VOQUEZNA DUAL PAK?

VOQUEZNA TRIPLE PAK or VOQUEZNA DUAL PAK can cause serious side effects including:

- **Rash in patients with mononucleosis:** Amoxicillin (a component of VOQUEZNA TRIPLE PAK and VOQUEZNA DUAL PAK) may cause a rash in patients who have mononucleosis. Stop taking VOQUEZNA TRIPLE PAK or VOQUEZNA DUAL PAK if you are diagnosed with mononucleosis and call your doctor right away.
- **Altered test results for some tumors:** VOQUEZNA TRIPLE PAK and VOQUEZNA DUAL PAK lower stomach acid which can cause increased levels of a certain protein (CgA) in your blood. When this level is increased it may alter test results for detecting some tumors. Notify your doctor of the use of VOQUEZNA TRIPLE PAK or VOQUEZNA DUAL PAK prior to blood tests.



Take your medication 3 times per day: in the morning, mid-day, and evening.



Pills not actual size.

**14-day treatment** that can be taken with or without food

### IF YOU MISS A DOSE OF VOQUEZNA® DUAL PAK®

Take the next dose as soon as possible unless 4 hours have passed since you were supposed to take it. If more than 4 hours have passed, skip the missed dose and take the next dose on time. Continue to take your medicine until all doses in the pack have been taken, even if symptoms go away.

## IMPORTANT SAFETY INFORMATION (cont'd)

Additionally, VOQUEZNA TRIPLE PAK can cause:


- **Irregular heartbeats:** Clarithromycin may cause irregular heartbeats. Call your doctor right away if you feel faint, light headed, or feel your heart beating irregularly.
- **Liver problems:** Call your doctor right away if you have any of the following symptoms: weight loss, yellowing of the skin and eyes (jaundice), dark urine, rash, or pain on the right side of your abdomen.

The most common side effects may include:

- diarrhea
- temporary changes in sense of taste
- vaginal yeast infection
- stomach pain
- headache
- highblood pressure and
- cold-like symptoms

Please see additional Important Safety Information throughout and full [Prescribing Information](#).

## SAVINGS AND SUPPORT OFFERINGS



### Your physician may send your prescription at BlinkRx\*

BlinkRx helps identify potential savings available to you. Phathom Pharmaceuticals and BlinkRx work together to help make it simple for you to receive VOQUEZNA® Triple Pak® or VOQUEZNA® Dual Pak®.



**Easy to use.** BlinkRx coordinates with your insurance on your behalf.



**Get the lowest out-of-pocket cost available to you.** BlinkRx identifies the cost of your medication and what insurance coverage you have, and then applies any savings offers for which you are eligible.



**Free home delivery.** A pharmacy partner will deliver your Rx free to your door.

After your doctor sends in your prescription, BlinkRx will contact you with details about how to register and use their service. You also have the option to register yourself now at 1 (844)-759-0782.

### You may be eligible for this savings card

If you have commercial insurance, you may be able to save on your prescription. You can sign up for a savings card [here](#).



For full Terms and Conditions click [here](#).

### IMPORTANT SAFETY INFORMATION (cont'd)

These are not all of the possible side effects of VOQUEZNA TRIPLE PAK and VOQUEZNA DUAL PAK. Call your healthcare provider for medical advice about side effects.

\*BlinkRx is NOT insurance. By sending your prescription to BlinkRx, you authorize BlinkRx ("Blink") to send text message notifications related to your prescriptions. Message and data rates may apply. Message frequency varies. Standard SMS messaging may not be fully secure. Providing consent is not a condition of purchasing goods or services through Blink. You can reply to any text with HELP for help or STOP to stop and revoke your consent. Visit Blink's HIPAA Privacy Notice and Terms of Use at [www.blinkrx.com/legal](http://www.blinkrx.com/legal).

Please see additional Important Safety Information throughout and full [Prescribing Information](#).



## FDA-APPROVED: A UNIQUE 14-DAY TREATMENT



Contains the acid blocker vonoprazan—which begins blocking acid quickly and helps improve the ability of antibiotics to kill *H. pylori* bacteria



Was proven in a clinical trial to get rid of *H. pylori* in most patients after 14 days of treatment



Can be taken with or without food



The most common side effects may include: diarrhea, temporary changes in sense of taste, vaginal yeast infection, stomach pain, headache, high blood pressure, and cold-like symptoms

**For more information, contact your doctor or visit [VOQUEZNA.com/HP](https://www.voquezna.com/HP)**

## IMPORTANT SAFETY INFORMATION (cont'd)

### General information about the safe and effective use of VOQUEZNA.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use VOQUEZNA TRIPLE PAK or VOQUEZNA DUAL PAK for a condition for which it was not prescribed. Do not give VOQUEZNA TRIPLE PAK or VOQUEZNA DUAL PAK to other people, even if they have the same symptoms you have. It may harm them.

For more information, ask your healthcare provider or pharmacist.

**You are encouraged to report suspected adverse reactions by contacting Phathom Pharmaceuticals at 1-888-775-PHAT (7428 or FDA at 1-800 FDA-1088 or [www.fda.gov/medwatch](https://www.fda.gov/medwatch).**

**Please see additional Important Safety Information throughout and full [Prescribing Information](#).**