

Narasin and Nicarbazin

Type B Medicated Broiler Chicken Feed

Do Not Feed Undiluted.

IMPORTANT: MUST BE THOROUGHLY MIXED INTO FEED BEFORE USE

INDICATIONS

For the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

ACTIVE DRUG INGREDIENTS

Narasin ³	46 g/ton to 11,350 g/ton*
Nicarbazin ³	46 g/ton to 11,350 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Methionine, not less than	_____	%
Crude Fat, not less than	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ¹ , not less than.....	_____	%
Salt ¹ , not more than.....	_____	%
Sodium ² , not less than	_____	%
Sodium ² , not more than	_____	%

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

MIXING DIRECTIONS

Thoroughly mix narasin and nicarbazin Type B Medicated Feed into complete broiler chicken feed to provide 27 – 45 g/ton narasin and 27 – 45 g/ton nicarbazin in the Type C Medicated Feed (Table 1).

Table 1: Mixing Directions for Broiler Chicken Medicated Feed

Narasin and Nicarbazin Concentration in Type B Feed (g/ton)	Desired Narasin and Nicarbazin Concentration in Type C Medicated Feed (g/ton)**	
	27 and 27	45 and 45
	lb of Type B per ton of Narasin and Nicarbazin Medicated Complete Feed	
46 and 46	1174	1956
5,675 and 5,675	9.5	15.9
11,350 and 11,350	4.8	7.9

CAUTION: For broiler chickens only. Do not feed to laying hens. Do not allow adult turkeys, horses or other equines access to formulations containing narasin. Ingestion of narasin by these animals has been fatal.

WARNING:

Withdraw 5 days before slaughter.

MANUFACTURED BY
BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk

*The medicated feed label must state a single drug concentration.

**The two drugs can be combined only at a 1:1 ratio for the 27 to 45 g/ton range.

³Sourced from Maxiban, NADA#138-952.



Elanco, Maxiban and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

YL090146AMA