

Tylosin Liquid Cattle Feed

Type B Medicated Feed

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Do Not Feed Undiluted

INDICATION

For reduction of incidence of liver abscesses associated with *Fusobacterium necrophorum* and *Arcanobacterium pyogenes*.

ACTIVE DRUG INGREDIENT

Tylosin phosphate^a 11 to 20,000 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Non-Protein Nitrogen (NPN) ¹ , not more than	_____	%
Crude Fat, not less than	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ² , not less than.....	_____	%
Salt ² , not more than.....	_____	%
Sodium ³ , not less than	_____	%
Sodium ³ , not more than	_____	%
Potassium, not less than	_____	%
Vitamin A ^{2,4} , not less than.....	_____	I.U./lb
Dry Matter, not less than	60%	
Dry Matter, not more than	75%	
pH.....	4.5 to 6.0	

¹ When added.

² If added.

³ Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴ Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

MIXING DIRECTIONS

Maintain the pH between 4.5 and 6.0.

Thoroughly mix Tylan Type B Medicated Feed into one ton of complete cattle feed to obtain the correct concentration in the Type C Medicated Feed (Table 1).

For liquid feeds stored in recirculating tank systems: Recirculate immediately prior to use for not less than 10 minutes, moving not less than 1 percent of the tank contents per minute from the bottom of the tank to the top. Recirculate daily as described even when not used.

For liquid feeds stored in mechanical, air or other agitation-type tank systems: Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily as described even when not used.

Table 1: Mixing Directions for Cattle Medicated Feed

Tylosin Concentration in Type B Feed (g/ton)	Desired Tylosin Concentration in Type C Medicated Feed (g/ton)	
	8	10
	lb of Type B per ton of Tylosin Medicated Complete Feed	
200	80	100
2,000	8	10
20,000	0.8	1

CAUTION

Do not use in any liquid feed containing sodium metabisulfite or in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

Expiration date: 31 days after the date of manufacture

MANUFACTURED BY:

BLUE BIRD FEED MILL

Any Town USA, 12345

Net Weight lb (kg) on bag or bulk

^a Sourced from Tylan, NADA# 12-491

*The medicated feed label must state a single drug concentration.

Elanco, Tylan and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

