# Tylosin Medicated Feed - Swine Dysentery with Tylosin in Drinking Water Type B Medicated Feed

**CAUTION:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

### **Do Not Feed Undiluted**

#### **INDICATION**

For treatment and control of swine dysentery associated with Brachyspira hyodysenteriae immediately after medicating with tylosin in drinking water.

#### **ACTIVE DRUG INGREDIENT**

101 to 20,000 g/ton*
%
%
%
%
%
%
%
%
%
%
%
ppm
ppm

<sup>1</sup> If added.

#### INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

#### MIXING DIRECTIONS

Thoroughly mix Tylosin Type B Medicated Feed into one ton of complete swine feed to obtain the correct concentration in the Type C Medicated Feed (Table 1).

Table 1: Mixing Directions for Swine Medicated Feed

Tylosin Concentration	Desired Tylosin Concentration in Type C Medicated Feed (g/ton)		
in Type B Feed (g/ton)	40	100	
	Ib of Type B per ton of Tylosin Medicated Complete Feed		
200	400	1000	
2,000	40	100	
20,000	4	10	

## **CAUTION**

Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

## **MANUFACTURED BY:**

**BLUE BIRD FEED MILL** Any town, USA 12345

## Net Weight lb (kg) on bag or bulk



<sup>&</sup>lt;sup>2</sup> Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>&</sup>lt;sup>a</sup> Sourced from Tvlan, NADA# 12-491

<sup>\*</sup>The medicated feed label must state a single drug concentration.