

BLUE BIRD TURKEY FEED

TYPE C MEDICATED FEED

INDICATIONS

For the prevention of coccidiosis caused by *E. adenoeides*, *E. meleagrimitis*, and *E. gallopavonis*, and as an aid in the control for transmissible enteritis in growing turkeys complicated by organisms susceptible to bacitracin methylene disalicylate.

ACTIVE INGREDIENT

Monensin¹ (as monensin sodium) 54 to 90 g/ton*
Bacitracin Methylene Disalicylate.....200 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than _____ %
Crude Fat, not less than _____ %
Crude Fiber, not more than _____ %

INGREDIENTS

Each ingredient must be specifically named (unless stated as such in the guaranteed analysis listing) in accordance with the names and definitions listed in Title 21 CFR 501.110.

DIRECTIONS FOR USE

Feeding Directions: Feed continuously as the sole ration. The optimum level depends upon the severity of coccidiosis.

CAUTION: Do not allow horses, other equines, mature turkeys or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. Some strains of turkey coccidia may be monensin tolerant or resistant.

Monensin may interfere with the development of immunity to turkey coccidiosis.

Store in a cool, dry place.

MANUFACTURED BY

BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk

Lot No. _____

*The medicated feed label must state a single drug concentration.

¹Sourced from Coban, NADA#038-878

