

Net Weight on Bag or Bulk

BLUE BIRD BROILER RATION

TYPE C MEDICATED FEED

INDICATIONS

For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, and for increased rate of weight gain and improved feed efficiency in broiler chickens.

ACTIVE DRUG INGREDIENTS

Narasin ³	27 to 45 g/ton
Nicarbazin ³	27 to 45 g/ton
Bacitracin Methylene Disalicylate.....	4 to 50 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Methionine, not less than	_____	%
Crude Fat, not less than	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than	_____	%
Phosphorus, not less than.....	_____	%
Salt ¹ , not less than.....	_____	%
Salt ¹ , not more than.....	_____	%
Sodium ² , not less than	_____	%
Sodium ² , not more than	_____	%

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials' Publication (AAFCO Official Publication).

FEEDING DIRECTIONS

Feed continuously as the sole ration.

WARNING: Withdraw 5 days before slaughter.

CAUTION: For broiler chickens only. Do not feed to laying hens. Do not allow adult turkeys, horses or other equines access to formulations containing narasin. Ingestion of narasin by these species has been fatal. Nicarbazine medicated broilers may show reduced heat tolerance of exposed to high temperature and high humidity. Provide adequate drinking water and ventilation during these periods

MANUFACTURED BY

Blue Bird Feed Mills
Anytown, USA 12345

**The medicated feed label must state a single drug concentration.

³Sourced from Maxiban, NADA#138-952.

