

AviPro®

108 FC3 Platinum

Description: *Pasteurella multocida* bacterin, (Types 1, 4 and 3 & 4), Oil Emulsion.

Directions for use: Vaccine should be warmed to room temperature before use. Shake well before and during use. Inject subcutaneously or intramuscularly using aseptic technique. Use entire contents of bottle when first opened. Vaccinate chickens between 12 and 16 weeks of age. Vaccinate turkeys between 20 and 24 weeks of age. Revaccinate 4 to 6 weeks later. Revaccinate during molt.

Dose: ¼ ml (0.25 ml) per bird. Inject all birds in flock.

Recommended for the vaccination of chickens and turkeys as an aid in the prevention of Fowl Cholera, caused by *Pasteurella multocida*, Type 1 in chickens and Types 4 and 3X4 in turkeys. If fowl cholera is diagnosed before it has spread throughout the flock, it is often possible to check its spread by immediate vaccination of healthy birds.

Caution: Store at 2° –7°C (35° –45°F). Do not freeze.

Warning: Do not vaccinate within 42 days before slaughter. In case of accidental self-injection, seek medical attention immediately. Contact Elanco with questions or concerns at 1-800-428-4441 or 1-888-545-5973.

LIST NO: KV:8816-H

FOR VETERINARY USE ONLY
Further information: www.elanco.us

The label contains complete use information, including cautions and warnings. Always read, understand and follow the label and use directions.

Elanco®, AviPro®, Full Value Poultry™ and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries, or affiliates.

© 2016 Eli Lilly and Company, its subsidiaries or affiliates.
USPBUAVP00027



FULL VALUE POULTRY™

Elanco™

1-800-428-4441
www.elanco.us