

Monensin Medicated Calf Feed Type C Medicated Feed For Calves (excluding veal calves)

INDICATIONS

For the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

Active Drug Ingredient

Monensin, USP⁵ 10 to 200 g/ton*

Guaranteed Analysis

Crude Protein, not less than %
 Non-Protein Nitrogen (NPN)¹, not more than %
 Crude Fat, not less than %
 Crude Fiber, not more than %
 Calcium, not less than %
 Calcium, not more than %
 Phosphorus, not less than %
 Salt², not less than %
 Salt², not more than %
 Sodium³, not less than %
 Sodium³, not more than %
 Potassium, not less than %
 Vitamin A^{2,4}, not less than I.U./lb

¹When added.

²If added.

³Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴Other than precursors of Vitamin A.

Ingredients

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

*Final printed label on formulated Type C medicated feed must bear a single drug concentration.

Feeding Directions

Hand feed continuously to calves (excluding veal calves) to provide 0.14 to 1.00 mg monensin/lb. bodyweight, depending upon severity of challenge, up to a maximum of 200 mg monensin/hd/day.

Example for 30 g monensin per ton complete feed:

Calf Weight (lb)	Feed per day (lb)		Monensin intake (mg/day)	
	Min	Max	Min	Max
100	0.93	<i>ad libitum</i>	14	60**
200	1.86	<i>ad libitum</i>	28	120**
300	2.80	<i>ad libitum</i>	42	180**
400	3.73	13.3	56	200

**Limited by appetite, assumed to be not more than 4% of body weight

Example for 200 g monensin per ton supplement:

Calf Weight (lb)	Feed per day (lb)		Monensin intake (mg/day)	
	Min	Max	Min	Max
100	0.14	1	14	100
200	0.28	2	28	200
300	0.42	2	42	200
400	0.56	2	56	200

Caution

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Warning

A withdrawal period has not been established for monensin in pre-ruminating calves. Do not use in calves to be processed for veal.

Manufactured By

Blue Bird Feed Mill

Anytown, USA 12345

Net Weight lb on bag or bulk

⁵Sourced from Rumensin, NADA#095-735

