

Monensin Medicated Quail Feed

(monensin Type C medicated feed)

INDICATION

For the prevention of coccidiosis in growing Bobwhite quail caused by *Eimeria dispersa* and *E. lettyae*.

ACTIVE DRUG INGREDIENT

Monensin^a, USP 73 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Methionine, not less than	_____	%
Crude Fat, not less than	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than	_____	%
Phosphorus, not less than.....	_____	%
Salt ¹ , not less than.....	_____	%
Salt ¹ , not more than.....	_____	%
Sodium ² , not less than	_____	%
Sodium ² , not more than	_____	%

¹ If added.

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed 73 g/ton of complete feed. Feed continuously as the sole ration.

WARNING

Do not feed to laying chickens. Do not feed to chickens over 16 weeks of age.

CAUTION

Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. Not for replacement chickens intended to become broiler breeding chickens.

MANUFACTURED BY

BLUE BIRD FEED MILL

Any town, USA 12345

Net Weight lb (kg) on bag or bulk

^a Sourced from Coban, NADA# 38-878 and 130-736

Elanco, Coban and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

