

Pediatric Dosing Chart

Dosage for infants and children from your healthcare professional

DOSE: Every 6-8 hours as needed. Use only the dosing device (dosing syringe or dosing cup) that came with the product.

DO NOT use any other dosing device.

DO NOT give more than 4 doses in 24 hours. If possible, use weight to dose; otherwise, use age.

Use products **ONLY** as directed. Always read and follow the label on all MOTRIN® products.

Infants' MOTRIN® Concentrated Drops

Available in: **Berry, Dye-Free Berry**



WEIGHT	12-17 lbs	18-23 lbs
AGE	6-11 mos	12-23 mos
DOSE	1.25 mL	1.875 mL



Children's MOTRIN® Oral Suspension

Available in: **Grape, Berry, Dye-Free Berry, Bubblegum**



WEIGHT	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs
AGE	2-3 yrs	4-5 yrs	6-8 yrs	9-10 yrs	11 yrs
DOSE	5 mL	7.5 mL	10 mL	12.5 mL	15 mL




Children's MOTRIN® Chewables

Available in: **Grape, Dye-Free Grape**



WEIGHT	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs
AGE	2-3 yrs	4-5 yrs	6-8 yrs	9-10 yrs	11 yrs
DOSE	1 tab	1.5 tabs	2 tabs	2.5 tabs	3 tabs



Today's date: _____

This dosing recommendation from your healthcare professional will expire in 14 DAYS.

ACTIVE INGREDIENT: Children's MOTRIN® Oral Suspension and Chewables products have 100 mg of ibuprofen (NSAID)* in each 5mL/tablet. Infants' Concentrated Drops has 50 mg of ibuprofen (NSAID)** in each 1.25 mL.

*Product eligibility may vary by plan. Check with your benefits administrator to determine which expenses are eligible before you shop. **Nonsteroidal anti-inflammatory drug. Use products only as directed. © Kenvue Brands LLC 2025 · CTCM-79161