Pediatric Dosing Chart

Dosage for children from your healthcare professional

Use **ONLY** with dosing cup provided with product. **DO NOT** use any other dosing device.

ASK A DOCTOR before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. **DO NOT USE** if you've had an allergic reaction to ZYRTEC® products or ingredients, or to an antihistamine containing hydroxyzine. Use products ONLY as directed. Always read and follow the label on all ZYRTEC® products.

Children's ZYRTEC® Allergy Syrup*





AGE	under 2 yrs	2 to under 6 yrs	6 yrs and over
DOSE	Ask a doctor	2.5 mL or 5 mL	5 mL or 10 mL
		2.5 or 5 mL	5 or 10 mL
		Max dose 5 mL once daily or 2.5 mL every 12 hours.	Max dose 10 mL once daily or 5 mL every 12 hours.

Children's ZYRTEC® **Dissolve Tabs***






AGE	under 6 yrs	6 yrs and over	
DOSE	Ask a doctor	1 tab	One tab (10 mg) once daily.

A 5mg product may be appropriate for less severe symptoms.

Children's ZYRTEC® Chewables 2.5mg*





AGE	under 2 yrs	2 to 6 yrs	
DOSE	Ask a doctor	1 or 2 tabs	Max dose 2 tabs (5 mg) once daily or 1 tab (2.5 mg) every 12 hours.

Chew or crush tablets completely before swallowing.

Today's date: _

This dosing recommendation from your healthcare professional will expire in 14 DAYS.

Children's ZYRTEC® Chewables 10mg*

Available in: Grape



A	GE	under 6 yrs	over 6 yrs	
D	OSE	Ask a doctor	1 tab	One tab (10 mg) once daily.

A 5mg product may be appropriate for less severe symptoms. Chew or crush tablets completely before swallowing.

ACTIVE INGREDIENT: Cetirizine

*Relieves sneezing; runny nose; itchy, watery eyes; and itching of the nose or throat. © Johnson & Johnson Consumer Inc. 2024 · ZYR-61037

























