

Pediatric Dosing Chart



Dosage for children from your healthcare professional

Ask a doctor or pharmacist before use if your child is presently taking a prescription drug. Antacids may interact with certain drugs. **DO NOT** use any other dosing device. Use products **ONLY** as directed. Always read and follow the label on all MYLICON® products.

Children's MYLICON® Antacid Liquid



Available in: **Bubblegum**

AGE	under 2 yrs	2 to 5 yrs	6 to 11 yrs
WEIGHT	under 24 lbs.	24 - 47 lbs.	48 - 95 lbs.
DOSE	Ask a doctor	5 mL 	10 mL 

DOSING: 5 mL - not more than 15 mL in 24 hours.
10 mL - not more than 30 mL in 24 hours.

Infants' MYLICON® Gas Relief Drops



Available in: **Original, Dye Free**



AGE	under 2 yrs	over 2 yrs
WEIGHT	under 24 lbs.	over 24 lbs.
DOSE	0.3 mL	0.6 mL

Can be used at every feeding, up to 12 times a day.

Children's MYLICON® Antacid Chewables



Available in: **Cherry**

AGE	under 2 yrs	2 to 5 yrs	6 to 11 yrs
WEIGHT	under 24 lbs.	24 - 47 lbs.	48 - 95 lbs.
DOSE	Ask a doctor	1 tab 	2 tabs 

DOSING: 1 tablet - not more than 3 tablets in 24 hours.
2 tablets - not more than 6 tablets in 24 hours.

Infants' MYLICON® Liquid Probiotic



Available in: **Newborns+**

AGE	Newborns
WEIGHT	—
DOSE	6 drops a day

DOSING: Do not exceed 6 drops in 24 hours.

Today's date: _____

This dosing recommendation from your healthcare professional will expire in 14 DAYS.

© Johnson & Johnson Consumer Inc. 2024 · MYC-1134

ACTIVE INGREDIENTS: Calcium carbonate, simethicone (Children's MYLICON® Antacid Liquid, Children's MYLICON® Antacid Chewables,) **Simethicone** (Infants' MYLICON® Gas Relief Drops,) **Bifidobacterium animalis subsp. lactis** (Infants' MYLICON® Liquid Probiotic.)