

# Pediatric Dosing Chart

Dosage for children from your healthcare professional

**DO NOT USE** with any other product containing diphenhydramine, even one used on skin.

**DO NOT** use any other dosing device.

**DO NOT** use to make a child sleepy.

Use products **ONLY** as directed. Always read and follow the label on all BENADRYL® products.

## Children's BENADRYL® Allergy Liquid\*

Available in: **Cherry, Bubblegum**



| AGE  | under 2 yrs | 2 to 5 yrs   | 6 to 11 yrs |
|------|-------------|--------------|-------------|
| DOSE | Do not use  | Ask a doctor | 5 or 10 mL  |

5 mL or 10 mL

Dose every 4 to 6 hours.  
DO NOT give more than 6 doses in 24 hours.

## Children's BENADRYL® Allergy Plus Congestion\*

Available in: **Grape**



| AGE  | under 4 yrs | 4 to 5 yrs   | 6 to 11 yrs | over 12 yrs |
|------|-------------|--------------|-------------|-------------|
| DOSE | Do not use  | Ask a doctor | 5 mL        | 10 mL       |

5 mL      10 mL

Dose every 4 hours.  
DO NOT give more than 6 doses in 24 hours.

## Children's BENADRYL® Chewables\*

Available in: **Grape**



| AGE  | under 2 yrs | 2 to 5 yrs   | 6 to 11 yrs | 12 yrs and over |
|------|-------------|--------------|-------------|-----------------|
| DOSE | Do not use  | Ask a doctor | 1 or 2 tabs | 2 or 4 tabs     |

or

Dose every 4 to 6 hours.  
DO NOT give more than 6 doses in 24 hours.

Today's date: \_\_\_\_\_

This dosing recommendation from your healthcare professional will expire in 14 DAYS.

**ACTIVE INGREDIENT: Diphenhydramine (Antihistamine)**

\*Relieves runny nose, sneezing, itchy, watery eyes; and itching of the nose or throat.

© Johnson & Johnson Consumer Inc. 2024 · ZYR-61037