

Sports Injuries

Whether they're playing sports or just a simple game of tag, kids can sometimes overdo it. Help them feel better and stay safe.



Play smart

- Keep sports fun and age-appropriate
- Help skills and muscles develop through different kinds of activities
- Limit sports to 5 days a week and 1 sport per season
- Schedule a physical exam for your child 4 to 6 weeks before the season starts



Know the signs

"Overuse injuries"
happen when bones,
muscles, and other
body parts are used too much
without time to heal. Talk to a
healthcare professional right
away if your child:

- · Can't put weight on a body part
- · Favors one side of the body
- Says it hurts to use a body part
- · Can't sit and/or climb stairs
- · Has stiff joints or muscles
- · Is unusually weak
- · Has trouble breathing
- Has dizziness, lightheadedness, or headache
- Loses feeling in fingers or toes

Remember R.I.C.E.

For strains and sprains, follow these steps:



- Rest your child from their regular activities and have them keep weight off the injury
- **Ice** the injured area for ONLY 20 minutes, 4 to 8 times daily
- Compress the injured area with an elastic wrap, a boot, or a splint to reduce swelling
- Elevate the injured area above the heart to further reduce swelling

Tackle pain and fever



Т)	YLI	ENO)L@

Active ingredient: acetaminophen

- Starts to reduce fever quickly
- Reduces pain and fever while being gentle on tummies

MOTRIN®

Active ingredient: ibuprofen

- Treats pain and fever
- Works fast and lasts up to 8 hours

Rely on **TYLENOL**° and **MOTRIN**°, the two brands of pain and fever relievers that pediatricians recommend most.*

Health tips from the American Academy of Pediatrics (AAP) and the Centers for Disease Control and Prevention (CDC)



Dosage On Call

Dosage for infants and children from your healthcare professional

DOSE: Every 4 hours as needed. Do not give more than 5 doses in 24 hours.

If possible, use weight to dose; otherwise, use age.

Do **NOT** use with any other product containing acetaminophen.

DOSE: Every 6-8 hours as needed. Do not give more than 4 doses in 24 hours.

If possible, use weight to dose; otherwise, use age.

 \mathbf{mL} = milliliter



Infants' TYLENOL® Oral Suspension

Active ingredient: acetaminophen 160 mg (in each 5 mL)

Available in: PGrape Cherry a Dye-Free Cherry							
WEIGHT	6-11 lbs	12-17 lbs 18-23 lbs 24-35 lbs					
AGE	0-3 months	4-11 months	12-23 months	2-3 years			
DOSE	1.25 mL	2.5 mL	3.75 mL	5 mL			
	2- 25 2: 25	2- 25 2 25	2-2522	2- 25 25 25			



Infants' MOTRIN® Concentrated Drops

Active ingredient: ibuprofen (NSAID)* 50 mg (in each 1.25 mL)

Available in: Berry Dye-Free Berry					
WEIGHT	12-17 lbs	18-23 lbs			
AGE	6-11 months	12-23 months			
DOSE	1.25 mL	1.875 mL			
	A S A S A S	2 % 2 % 2 % A % A % A % A % A % A % A %			



product

Children's TYLENOL® Oral Suspension

Active ingredient: acetaminophen 160 mg (in each 5 mL)

Available	e in: 🦃 Grape	Cherry 🧟	Dye-Free Cherry	Bubblegum	Strawberry
WEIGHT	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs
AGE	2-3 years	4-5 years	6-8 years	9-10 years	11 years
DOSE	5 mL	7.5 mL	10 mL	12.5 mL	15 mL
	5 mL	7.5 mL	10 mL	12.5 mL	15 mL



Children's MOTRIN® Oral Suspension

Active ingredient: ibuprofen (NSAID)* 100 mg (in each 5 mL)

Available in: Frape Frape Serry Berry Bubble Gum					
WEIGHT	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs
AGE	2-3 years	4-5 years	6-8 years	9-10 years	11 years
DOSE	5 mL	7.5 mL	10 mL	12.5 mL	15 mL



Children's TYLENOL® Chewables

Active ingredient: acetaminophen 160 mg (in each chewable tablet)

Availabl	Available in: Bubblegum						
WEIGHT	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs		
AGE	2-3 years	4-5 years	6-8 years	9-10 years	11 years		
DOSE	1 tablet	1½ tablets	2 tablets	21/2 tablets	3 tablets		



only as directed.

only as directed

Children's MOTRIN® Chewables

Available in: Free Grape

 $\textbf{Active ingredient:} \ \text{ibuprofen (NSAID)*} \ 100 \ \text{mg (in each chewable tablet)}$

	_	_			
WEIGHT	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs
AGE	2-3 years	4-5 years	6-8 years	9-10 years	11 years
DOSE	1 tablet	11/2 tablets	2 tablets	21/2 tablets	3 tablets



Children's TYLENOL® Dissolve Packs

Active ingredient: acetaminophen 160 mg (in each pack)

Available in: Wild Berry

Use product only as directed.

WEIGHT	under 48 lbs	48-59 lbs	60-71 lbs	72-95 lbs
AGE	under 6 years	6-8 years	9-10 years	11 years
DOSE	do not use	2 powders	2 powders	3 powders

All Infants' TYLENOL® and Children's TYLENOL® products have the same strength of acetaminophen: 160 mg (in each 5 mL, tablet, or pack).

*Nonsteroidal anti-inflammatory drug.

IMPORTANT INSTRUCTIONS FOR PROPER USE

Today's date:

This dosing recommendation from your healthcare professional will expire in **14 DAYS**.

- Always read and follow the label on all TYLENOL® and MOTRIN® products.
- Use only the dosing device (dosing syringe or dosing cup)
 that came with the product. Do not use any other dosing device.