



DOSING FOR HEALTHCARE PROFESSIONALS

Use this chart to determine the proper dose of **MOTRIN®** for your patient.
If possible, use weight to dose; otherwise, use age.†

† For pediatric products only

Recommended dose of **MOTRIN®**
from your healthcare professional:

_____ dose of _____ product every _____ hours for _____ days

Visit Motrin.com for additional product information		Dosage					
		Infants' MOTRIN® Concentrated Drops	Children's MOTRIN® Oral Suspension	Children's MOTRIN® Chewables	MOTRIN® IB Caplets	MOTRIN® IB Liquid Gels	MOTRIN® IB Migraine Liquid Gels
mL = milliliter							
Weight (lbs)	Age	Active ingredient: ibuprofen (NSAID)* 50 mg (in each 1.25 mL) <small>*Nonsteroidal anti-inflammatory drug.</small>	Active ingredient: ibuprofen (NSAID)* 100 mg (in each 5 mL) <small>*Nonsteroidal anti-inflammatory drug.</small>	Active ingredient: ibuprofen (NSAID)* 100 mg (in each chewable tablet) <small>*Nonsteroidal anti-inflammatory drug.</small>	Active ingredient: ibuprofen (NSAID)* 200 mg (in each caplet) <small>*Nonsteroidal anti-inflammatory drug.</small>	Active ingredient: ibuprofen (NSAID)* 200 mg (in each capsule) <small>*Nonsteroidal anti-inflammatory drug.</small>	Active ingredient: ibuprofen (NSAID)* 200 mg (in each capsule) <small>*Nonsteroidal anti-inflammatory drug.</small>
12-17 lbs	6-11 mos	1.25 mL	Infants' MOTRIN® and Children's MOTRIN® are different concentrations. The markings on the included dosing cup correspond to the OTC label doses listed below.	-	OTC label does not include dosing for children under 12 years old.	OTC label does not include dosing for children under 12 years old.	OTC label does not include dosing for children under 18 years old.
18-23 lbs	12-23 mos	1.875 mL		-			
24-35 lbs	2-3 years	Infants' MOTRIN® and Children's MOTRIN® are different concentrations. The OTC label does not include dosing directions for children ages 2 years and older.	5 mL	1 tablet			
36-47 lbs	4-5 years		7.5 mL	1½ tablets			
48-59 lbs	6-8 yrs		10 mL	2 tablets			
60-71 lbs	9-10 yrs		12.5 mL	2½ tablets			
72-95 lbs	11 yrs		15 mL	3 tablets			
-	-	If needed, repeat dose every 6-8 hours Do NOT use more than 4 times a day	If needed, repeat dose every 6-8 hours Do NOT use more than 4 times a day	If needed, repeat dose every 6-8 hours Do NOT use more than 4 times a day	OTC LABEL STATES: Adults and children 12 years and older: Take 1 caplet every 4 to 6 hours while symptoms persist; if pain or fever does not respond to 1 caplet, 2 caplets may be used Children under 12 years: Ask a doctor Do NOT take more than 6 caplets in 24 hours unless directed by a doctor	OTC LABEL STATES: Adults and children 12 years and older: Take 1 capsule every 4 to 6 hours while symptoms persist; if pain or fever does not respond to 1 capsule, 2 capsules may be used Children under 12 years Ask a doctor Do NOT take more than 6 capsules in 24 hours unless directed by a doctor	OTC LABEL STATES: Adults: Take 2 capsules with a glass of water; if symptoms persist or worsen, ask your healthcare professional Under 18 years of age: Ask a doctor Do NOT take more than 2 capsules in 24 hours unless directed by a doctor

Remind parents & caregivers:

- Read & follow the label on all **MOTRIN®** products.
- If needed, repeat dose every 6-8 hours while symptoms last.
- Do **NOT** give more than 4 doses in 24 hours.
- Use only the dosing device (syringe or dosing cup) that came with the product.
- *Nonsteroidal anti-inflammatory drug

Use product only as directed.
© Johnson & Johnson Consumer Inc. 2023

Important Instructions For Proper Use:

- Always read and follow the label on all **MOTRIN®** products. Keep carton.
- Wait the right amount of time between doses
- If you have any questions, ask your healthcare professional which product may be right for you
- *Nonsteroidal anti-inflammatory drug