MOTRIN® Dosage Made Easy

Strong, effective relief for minor aches and pains, including backaches, muscular aches, and minor arthritis pain



MOTRIN® IB Caplets

ACTIVE INGREDIENT	DOSE & FREQUENCY	DIRECTIONS
Ibuprofen (NSAID*) 200 mg (in each caplet)	Adults and children 12 years and over: Take 1 caplet every 4 to 6 hours while symptoms persist; if pain or fever does not respond to 1 caplet, 2 caplets may be used Children under 12 years: Ask a doctor	Do not take more than 6 caplets in 24 hours unless directed by a doctor

Fast-acting relief for minor aches and pains, including menstrual cramps, toothaches, and muscular aches



MOTRIN® IB Liquid Gels

ACTIVE INGREDIENT	DOSE & FREQUENCY	DIRECTIONS	
Ibuprofen (NSAID*) 200 mg (in each capsule)	Adults and children 12 years and over: Take 1 capsule every 4 to 6 hours while symptoms last; if pain or fever does not respond to 1 capsule, 2 capsules may be used Children under 12 years: Ask a doctor	Do not take more than 6 capsules in 24 hours unless directed by a doctor	

Fast, effective relief for migraine pain and migraine-related symptoms, such as light sensitivity and nausea



MOTRIN® Migraine Liquid Gels

ACTIVE INGREDIENT	DOSE & FREQUENCY	DIRECTIONS	
Ibuprofen (NSAID*) 200 mg (in each capsule)	Adults: Take 2 capsules with a glass of water; if symptoms persist or worsen, ask your healthcare professional Under 18 years of age: Ask a doctor	Do not take more than 2 capsules in 24 hours unless directed by a doctor	

Relief for minor aches and pains, with an effective nighttime sleep aid



Use product only as directed

	MC	ו	ΚI	N®	РМ	Cap	olets

ACTIVE INGREDIENT	DOSE & FREQUENCY	DIRECTIONS	
Diphenhydramine citrate 38 mg Ibuprofen (NSAID*) 200 mg (in each caplet)	Adults and children 12 years and over: Take 2 caplets at bedtime	Do not take more than 2 caplets in 24 hours	

^{*}Nonsteroidal anti-inflammatory drug.

This is not a complete list of all MOTRIN® products.

IMPORTANT INSTRUCTIONS FOR PROPER USE:

- Always read and follow the label on all MOTRIN® products.
 Keep carton.
- Do NOT use with any other product containing an NSAID*
- Wait the right amount of time between doses
- If you have any questions, ask your healthcare professional which product may be right for you



Visit Motrin.com for additional product information