

- Confirm that you have a 12 mg Humatrope Cartridge to match the HumatroPen 12 mg. If it does not match DO NOT USE and contact your physician, pharmacist or nursing staff.
- Follow Section 2 ONLY to set up a new Cartridge before first use.
- Section 3 of this manual should be used for every injection.
- DO NOT SHARE your HumatroPen 12 mg or Pen Needles as this may risk transmission of infectious agents.
- The HumatroPen 12 mg is not recommended for use by blind or visually impaired individuals without the assistance of a sighted individual trained in its use.

## ABOUT PEN NEEDLES

pharmacist or nursing staff.

## What kinds of Needles can be used with the HumatroPen 12 mg?

- Pen Needles are not included. You may need a prescription to get the Needles from your pharmacist.
- Becton, Dickinson and Company (BD) Pen Needles are recommended for use with the HumatroPen 12 mg.
- Use of the Hidden Needle Cover with your HumatroPen is optional.
- When using the Hidden Needle Cover with your HumatroPen, you must use a Needle length of 5 mm or greater.
  Refer to the Hidden Needle Cover Instructions for use
- Ask your physician, pharmacist or nursing staff what Needle gauge and length is best for you.

## Must a new Needle be used for each injection?

- Yes, a new Needle must be used for each injection.
- Remove the Needle immediately after each injection. Use a new Needle for each injection. This will help minimize the risk of infection, prevent leakage of Humatrope, keep out air bubbles, and reduce Needle clogs.

## How do I throw away used Needles?

• Throw away used Needles as directed by your physician, pharmacist or nursing staff.

## CARE AND STORAGE FOR THE HUMATROPEN 12 MG

## <u>Care</u>

- Soiled parts can be cleaned with a damp cloth. DO NOT USE alcohol or other cleaning agents.
- •
- DO NOT APPLY oil or any other lubricant.

## **Storage**

- •
- Store HumatroPen 12 mg between -40°C to 70 °C without the Humatrope (somatropin) 12 mg Cartridge.
- Store the HumatroPen 12 mg with attached Humatrope Cartridge in the storage case in the refrigerator until the time of the next injection. DO NOT FREEZE.
- Refer to the Humatrope cartridge Patient Information Leaflet for complete storage instructions.
- Let the HumatroPen 12 mg with attached Humatrope Cartridge stand at room temperature for 10 minutes before injecting. Discomfort may be noticed at the injection site if Humatrope is injected cold.
- Daily room temperature exposure should not exceed 30 minutes.
- DO NOT STORE the Pen with the Needle attached.

## your physician, pharmacist or nursing staff. **STEP B - ATTACH THE CARTRIDGE** N Use the White Tip of the Cartridge to Push the White Tip of a reconstituted CHECK Cartridge into the Pen Body. Screw push the Screw back. the 12 mg Pen Body onto the Look at the Injection Button, the Cartridge until it is secure. ΝΟΤΕ Front Housing and cartridge label to confirm it is a 12 mg Pen. The screw may not be out when you get the Pen. NOTE If the Cartridge is not completely attached, the Screw may not move, and an incorrect dose may be given. **STEP C - ATTACH THE NEEDLE** • Pull off the Outer Cap and the Remove the Paper Tab from the end Push the Needle **straight** onto the of the Outer Cap. 12 mg Cartridge and screw on Inner Cap. clockwise until secure. • Keep the Outer Cap to remove the Needle after your injection. **STEP D - REMOVE AIR FROM NEW CARTRIDGE** 1.30 / -2.50 -0.20 1.40



- Point the Needle straight up.Push the Injection Button and hold for five seconds.
- Dial 0.10 mg and repeat these three steps until you see a stream of liquid.

## **REPLACEMENT and DISPOSAL OF PEN**

Do not use your pen for more than 3 years after the first use or past the use-by date on the carton, whichever comes first. The expired pen may be discarded in your household waste after you have removed the needle. Ask your healthcare professional about options to dispose of the pen properly. Record the date the Pen was first used here:  $__/ ___/$ . Record the Use-by date from your carton here:  $__/ ___/$ .

- The Pen must be set up before injecting the first dose from each new 12 mg Cartridge.
- Setting up the new Cartridge is important to remove large air bubbles that may be present after reconstitution (mixing).
- If a stream is not seen after several attempts, contact your physician, pharmacist or nursing staff or your local Lilly affiliate (see last page for contact details).

ΝΟΤΕ

Contact your physician, pharmacist or nursing staff if a new HumatroPen 12 mg is needed, or when the Pen has been used for 3 years.

**i** Please see the complete Humatrope *Patient Information Sheet*.

For additional information, contact your physician, pharmacist or nursing staff or your local Lilly affiliate (see last page for contact details) or visit our website at www.lilly.co.uk

## **STEP E - CONTINUE ON TO DAILY USE**

- DO NOT REPEAT Cartridge Setup before each dose.
- Leave the Cartridge attached and DO NOT REMOVE until the Cartridge is empty.
- Go to Section 3, Step 3, for instructions on how to inject the first dose.





Now that you have done the one-time New Cartridge Setup, follow Section 3 for all of your injections.



## **COMMONLY ASKED QUESTIONS**

## 1. Do I need to perform the New Cartridge Setup before every dose?

- No. The New Cartridge Setup is done only once for each Cartridge, just before a new Cartridge is used for the first time.
- The purpose of the setup is to make sure the HumatroPen 12 mg with the Humatrope 12 mg Cartridge are ready to use.
- If you repeat the New Cartridge Setup before each routine dose, you may run out of Humatrope early. The small amount of product used in the New Cartridge Setup will not affect the supply of Humatrope.

## 2. What should I do if the Cartridge Label and Pen do not match?

- DO NOT USE the Pen if the Cartridge strength on the Humatrope Cartridge label does not match the number on the Pen's Front Housing.
- Contact your physician, pharmacist or nursing staff for assistance or to obtain a replacement.

## 3. What should I do if the Humatrope is not clear after mixing?

- Be sure to gently invert the Pen up and down 10 times. DO NOT SHAKE. Then, let the Pen sit for at least three minutes. If the solution remains cloudy or has particles, gently invert the Pen up and down 10 more times. Let the Pen sit for five more minutes.
- If the solution remains cloudy or contains particles after reconstitution (mixing), DO NOT USE. Contact your physician, pharmacist or nursing staff for assistance.

## 4. Why are there air bubbles in the Cartridge?

- Air bubbles may remain in the Cartridge after reconstitution (mixing).
- If the Pen is stored with a Needle attached, air bubbles may form in the Cartridge. **DO NOT STORE** the Pen with a Needle attached.
- Perform the New Cartridge Setup to remove air bubbles from the Cartridge.
- A small air bubble is normal. It will not harm you nor affect the dose.

## 5. Why doesn't the Screw move out when there is no Cartridge attached to the Pen?

- The Screw may not move out when you push the Injection Button unless there is a Cartridge in the Pen. This • feature allows you to easily push the Screw into the Pen Body when replacing a Cartridge.
- Once a Cartridge is attached, the Screw will move out when the Injection Button is pushed.

## 6. What should I do if I can't attach the Cartridge to the Pen Body?

- Check that the Cartridge is not damaged or broken.
- Carefully line up the Cartridge with the Pen Body and screw together until secure. If the Cartridge and Pen cannot be screwed together contact your physician, pharmacist or nursing staff.

## 7. Why is it difficult to push the Injection Button when I try to inject the dose?

- The Needle may be clogged. Try attaching a new Needle.
- Pushing the Injection Button down quickly may make the Injection Button harder to push. Pushing the Injection Button more slowly may make it easier.

# **DAILY USE**

# **STEP 1 - CHECK THE PEN** Pull off the Pen Cap. Be sure to check the Cartridge:

- For 12 mg Cartridge Label For expiration date • Contents should be clear and free
- of particles

# NOTE

## **DO NOT USE the Pen if any part** of the Pen or Cartridge appears broken or damaged. Contact your physician, pharmacist or nursing staff.

## Check that the number on the Front Housing matches the Cartridge strength on the Cartridge label. If the Pen and Cartridge do not match, contact your physician, pharmacist or nursing staff.

Pen.

## **STEP 2 - ATTACH THE NEEDLE**



• Remove the Paper Tab from the

end of the Outer Cap.



12


Look at the Injection Button and the

Front Housing to confirm it is a 12 mg

CHECK

## Push the Needle **straight** onto the • Pull off the Outer Cap and the 12 mg Cartridge and screw on Inner Cap. clockwise until secure.

• Keep the Outer Cap to remove the Needle after the injection.



• **DO NOT REMOVE** this Cartridge from the Pen until the Cartridge is empty or needs to be replaced - this will help to avoid the possibility of an inaccurate dose.

- Using a larger diameter Needle, for example a 12.7mm Needle (diameter 0.33mm) will make it easier to push the Injection Button during injection. Ask your physician, pharmacist or nursing staff which Needle is best for vou
- The Injection Button may become harder to push if the inside of your Pen gets dirty with Humatrope, food, drink or other materials.

## 8. Why doesn't the Dose Knob go to zero when I inject the dose?

• This can happen if the Humatrope Cartridge does not have enough Humatrope left in it for the full dose. It is possible to set a dose larger than the amount of Humatrope left in the Cartridge. At the end of the injection, the number in the Dose Window should be 0.00. If it is not, this is the amount of Humatrope you DID NOT receive. Consult with your physician, pharmacist or nursing staff on how to handle a partial dose. Remove the Needle and empty Cartridge. For the next daily use attach a new Cartridge as shown in Section 2, Step A, and continue with New Cartridge Setup (Section 2).

## 9. Why do I see Humatrope leaking from the Needle after I have finished the injection?

- It is normal for a single drop to remain on the tip of the Needle after the injection is complete. If you see more than one drop:
  - You may not have received your full dose. DO NOT INJECT another dose. Consult with your physician, pharmacist or nursing staff for assistance.
  - To prevent this, for your next dose, firmly push and hold the Injection Button in and slowly count to five (see Section 3, Step 3).

## 10. How can I tell when the injection is complete?

- The injection is complete when:
  - You have slowly counted to five while you are still holding the Injection Button in and before you remove the Needle from the skin.

## AND

- 0.00 is in the center of the Dose Window.

## If you have any questions or problems with your HumatroPen 12 mg, contact local Lilly affiliate office (see below) or your healthcare professional for assistance.

Report all MEDICAL DEVICE COMPLAINTS or UNDESIRABLE SIDE EFFECTS including SUSPECTED SERIOUS INCIDENTS, to local Lilly affiliate office (see below).

You are also encouraged to report all UNDESIRABLE SIDE EFFECTS, including SUSPECTED SERIOUS INCIDENTS to

United Kingdom: Yellow Card Scheme, Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the **Google Play or Apple App Store** 

Australia: www.tga.gov.au/reporting-problems

Malta: ADR Reporting, Website: www.medicinesauthority.gov.mt/adrportal

South Africa: https://www.sahpra.org.za/Publications/Index/8

Local Lilly affiliate contact details: **United Kingdom** Eli Lilly and Company Limited

Tel: +44 (0) 1256 315000 Australia

Eli Lilly Australia Pty Ltd 112 Wharf Road West Ryde NSW 2114 Tel: 1800 454 559

Malta Charles de Giorgio Ltd. Tel: + 356 25600 500

Importer Lilly S.A. Avenida Industria 30 28108 Alcobendas Madrid, Spain

## Eli Lilly and Company Pharmaceutical Delivery Systems

Lilly Corporate Center Indianapolis IN 46285 USA

CE 0344

South Africa: Eli Lilly (S.A.) (Pty) Limited Golden Oak House (Building E), Ballyoaks Office Park, 35 Ballyclare Drive, Bryanston, 2191 Tel: +27(11) 510 9300

Humatrope<sup>®</sup> and HumatroPen<sup>®</sup> and H Block designs are trademarks of Eli Lilly and Company.

The HumatroPen<sup>®</sup> 12 mg meets the current dose accuracy and functional requirements of ISO 11608-1 and is for use with Humatrope<sup>®</sup> 12 mg Cartridges only.

Copyright © 2008, 2020 Eli Lilly and Company. All rights reserved.

Store the HumatroPen 12 mg properly. (See "Care and Storage for the HumatroPen 12 mg" in <b>Section 1</b> of this pen user manual for more information.) When it is time for your next dose, go to Section 3, and repeat Steps 1 -5.	3528 BJ UTRECHT The Netherlands	MD Medical device <sup>2</sup> Single patient – multiple use <sup>2</sup> <sup>1</sup> Glossary includes non-harmonized symbols used in the labeling. <sup>2</sup> Symbol used only on the carton. Revision date: April 2020