



# Swift<sup>®</sup> SC

1Le

F

## Fungicide

A strobilurin fungicide for the control of foliar diseases in winter wheat and winter and spring barley.

MAPP 11227

A suspension concentrate formulation containing 500 g/L (45.09% w/w) trifloxystrobin.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

Bayer CropScience Limited  
230 Cambridge Science Park  
Milton Road, Cambridge CB4 0WB  
Telephone: 01223 226500

For 24 hour emergency information contact **Bayer CropScience Limited**  
Telephone: 00800 1020 3333



### Safety Information

#### SWIFT SC

Contains 500 g/L (45.09 % w/w) trifloxystrobin



### Warning

Very toxic to aquatic life with long lasting effects.  
Wear protective gloves.

If skin irritation or rash occurs: Get medical advice/attention

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

Contains trifloxystrobin, 1,2-Benzisothiazolin-3-one and 5-Chloro-2-methyl-isothiazol-3-one/2-Methyl-isothiazol-3-one. May produce an allergic reaction.

### IMPORTANT INFORMATION FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crops:	Winter wheat, and winter and spring barley.
Maximum individual dose:	0.5 litres of product per hectare
Maximum number of treatments:	2 per crop
Latest time of application:	35 days before crop harvest
Other specific restrictions:	No more than two applications of products containing QoI fungicides must be applied to any cereal crop per year.

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

PROTECT FROM FROST



To access the **Safety Data Sheet** for this product scan the code or use the link below:

[www.cropscience.bayer.co.uk/swiftscsds](http://www.cropscience.bayer.co.uk/swiftscsds)  
or alternatively contact your supplier

GB79015984j rA7b

Bayer

## SAFETY PRECAUTIONS

### Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate. However engineering control may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

### WASH ALL PROTECTIVE CLOTHING AFTER USE

DO NOT BREATHE SPRAY.

WASH SPLASHES from skin or eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

### Consumer Protection

DO NOT HARVEST CROPS FOR HUMAN OR ANIMAL CONSUMPTION FOR AT LEAST 35 DAYS AFTER THE LAST APPLICATION.

### Environmental Protection

**DO NOT CONTAMINATE SURFACE WATERS OR DITCHES with chemical or used container.**

### Storage and Disposal

KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling, and dispose of safely.

## DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

### RESTRICTIONS

An interval of at least 35 days must elapse between the last application of Swift SC and harvesting the crop.

PROTECT FROM FROST

STORE IN A COOL DRY PLACE

### DISEASES CONTROLLED

When applied as recommended Swift SC controls the following diseases:

#### Winter Wheat

*Septoria (S. nodorum & S. tritici)*

Brown rust (moderate control)

#### Winter & Spring Barley

*Rhynchosporium*

Brown rust

Net blotch

Swift SC may be used on all commercial varieties of winter wheat and winter and spring barley.

Apply Swift SC at 0.5 L/ha in a minimum of 200 L water/ha. Use a **MEDIUM** spray quality (as defined by BCPC). A spray pressure of 2-3 bar is recommended. Apply a maximum of 2 applications of Swift SC per crop per year.

Apply Swift SC protectively, before disease is established in the crop. Further applications may be needed if disease attack is prolonged.

### RESISTANCE

To avoid build-up of resistance do not apply this or any other product containing QoI fungicides more than two times per crop per year.

Swift SC contains trifloxystrobin, a member of the QoI cross-resistance group. Swift SC should be used preventatively and not relied on for its curative potential. It should not be used where disease is already established. Use Swift SC as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control, including where appropriate other fungicides with a different mode of action. You must not apply more than two sprays of QoI containing products to any cereal crop.

Swift SC must always be used in mixture with another product, containing an active ingredient from a different cross resistance group, that is recommended for control of the same target disease applied at a rate that will give robust control. Where strains of fungi resistant to QoI's occur, Swift SC is likely to be ineffective and should not be used.

Strains of barley powdery mildew resistant to QoI's are common in the UK.

Limiting the number of sprays is an important factor in delaying the build-up of resistant pathogen populations. There is a significant risk of widespread resistance occurring in *Septoria tritici* populations in the UK. Failure to follow resistance management action may result in reduced levels of disease control. Appropriate disease risk assessment or decision support systems must be used when deciding whether to treat with Swift SC.

The possible development of resistance to Swift SC cannot be predicted. Should it occur Swift SC may not give satisfactory control.

## **MIXING AND SPRAYING**

Thoroughly shake the pack before use. Add half the volume of clean water to the spray tank. Turn on the agitation and add the recommended quantity of Swift SC. Continue agitation at all times, including topping up the spray tank and during spraying and any stoppages until the spray tank is empty. Wash out all spray equipment thoroughly with water immediately after use. Sprayers should be thoroughly cleaned before use, and filters and jets checked for damage and blockages. Boom height should be adjusted to ensure even coverage of the crop, particularly at later growth stages. The correct height is one at which the spray from alternate nozzles meets just above the crop. In dense crops, at later growth stages, higher water volumes should be used. Rinse container thoroughly, by using an integrated pressure-rinsing device or manually rinsing three times. Add washings to sprayer at time of filling, and dispose of safely.

© Swift is a Registered Trade Mark of Bayer  
© Bayer CropScience 2018



# Swift<sup>®</sup> SC

1Le

F

## Fungicide

A strobilurin fungicide for the control of foliar diseases in winter wheat and winter and spring barley.

MAPP 11227

A suspension concentrate formulation containing 500 g/L (45.09% w/w) trifloxystrobin.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

Bayer CropScience Limited  
230 Cambridge Science Park  
Milton Road, Cambridge CB4 0WB  
Telephone: 01223 226500

For 24 hour emergency information contact **Bayer CropScience Limited**  
Telephone: 00800 1020 3333



### Safety Information

#### SWIFT SC

Contains 500 g/L (45.09 % w/w) trifloxystrobin



### Warning

Very toxic to aquatic life with long lasting effects.  
Wear protective gloves.

If skin irritation or rash occurs : Get medical advice/attention

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

Contains trifloxystrobin, 1,2-Benzisothiazolin-3-one and 5-Chloro-2-methyl-isothiazol-3-one/2-Methyl-isothiazol-3-one. May produce an allergic reaction.

### IMPORTANT INFORMATION FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crops:	Winter wheat, and winter and spring barley.
Maximum individual dose:	0.5 litres of product per hectare
Maximum number of treatments:	2 per crop
Latest time of application:	35 days before crop harvest
Other specific restrictions:	No more than two applications of products containing QoI fungicides must be applied to any cereal crop per year.

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

PROTECT FROM FROST



To access the **Safety Data Sheet** for this product scan the code or use the link below:

[www.cropscience.bayer.co.uk/swiftscsds](http://www.cropscience.bayer.co.uk/swiftscsds)

or alternatively contact your supplier

GB79015984j rA7b

Bayer