

10Le

An agricultural/horticultural/industrial wetting agent for use with all authorised herbicides in maize (forage and grain), cereals, sugar beet, ornamental plant production, amenity vegetation, natural surfaces not intended to bear vegetation, permeable surfaces overlying soil, and industrial and amenity areas, and with insecticides in brassica crops and sugar beet.

ADJ number: 0818

An emulsifiable concentrate containing 81.4% w/w oil (rapeseed fatty acid esters) (EAC1).

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

Bayer CropScience Limited 230 Cambridge Science Park Milton Road Cambridge CB4 0WB Telephone: 01223 226500

For 24 hour emergency information contact Bayer CropScience Limited **Telephone: 00800 1020 3333**

MERO

UFI: E660-M0FM-P00N-FG11

Contains 81.4 % w/w oil (rapeseed fatty acid esters) (EAC1) and ethoxy (7) tridecanol.

Wear protective gloves/protective clothing/eye protection/face protection.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

Plant protection product	Crops		
All authorised herbicides	Forage and grain maize, cereals, sugar beet, ornamental plant production, amenity vegetation, natural surfaces not intended to bear vegetation, permeable surfaces overlying soil, and industrial and amenity areas		
All authorised insecticides	Brussels sprouts, cabbage, broccoli/calabrese, cauliflower and sugar beet		
Maximum concentration (% spray solution):	1.0		
Maximum number of treatments:	Follow the statutory conditions of use of the plant protection product.		
Latest time of application:	Follow the statutory conditions of use of the plant protection product, and: Maize: up to and including GS 53 Cereals: up to and including GS 52 Sugar beet: up to and including 6 leaves unfolded (GS 16) Brussels sprouts: up to and including lateral buds begin to develop (GS 41) Cabbage (Head): Up to heads begin to form (GS 41) Cabbage (leaf): Up to 10% of the leaf mass reached (GS 41) Cauliflower, broccoli/calabrese: Up to and including heads begin to form growing tip up to 1 cm ² (GS 41) Non-edible crops and non-crop production: follow the statutory conditions of use of the plant protection product		
Other specific restrictions:	Follow all operator protection instructions for the plant protection product. Where the personal protective equipment specified on the adjuvant label is different to that specified on the plant protection product label, the highest level of protective equipment must be worn.		



PRODUCTS.

To access the **Safety Data Sheet** for this product scan the code or use the link below:

www.cropscience.bayer.co.uk/merosds

or alternatively contact your supplier

Bayer CropScience Limited 230, Cambridge Science Park Milton Road, Cambridge CB4 0WB

Telephone: 01223 226500



Operator protection

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WASH CONCENTRATE from skin/eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

AVOID ALL CONTACT BY MOUTH.

WHEN USING DO NOT EAT DRINK OR SMOKE.

If swallowed, seek medical advice immediately and show this container or label.

Storage and Disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS. KEEP OUT OF REACH OF CHILDREN. KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place. WASH OUT CONTAINER THOROUGHLY and dispose of safely. DO NOT RE-USE CONTAINER FOR ANY OTHER PURPOSE.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the product label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

The maximum concentration of Mero in the spray solution is 1.0%. The quantity of Mero to add to different application volumes is as follows:

Application Volume (L/ha)	100	150	200	300
Required quantity of Mero (L)	1.0	1.5	2.0	3.0

Do not use Mero later than the crop growth stages listed below:

- Maize: up to and including tip of tassel visible (GS 53).
- Cereals: up to and including 1/4 inflorescence emerged (GS 52).
- Sugar beet: up to and including 6 leaves unfolded (GS 16).
- Brussels sprouts: up to and including lateral buds begin to develop (GS 41).
- Cabbage (head): Up to heads begin to form (GS 41).
- Cabbage (leaf): Up to 10% of the leaf mass reached (GS 41).
- Cauliflower, broccoli/calabrese: Up to and including heads begin to form, growing tip up to 1 cm² (GS 41).
- Non-edible crops and non-crop production: follow the statutory conditions of use of the plant protection product.

Herbicides

Weed control is not obtained by using Mero alone. When using Mero with an authorised herbicide, the statutory conditions of use of the plant protection product must be followed. Do not use Mero with an authorised herbicide later than GS 53 in maize, GS 52 in cereals or GS 16 in sugar beet.

Insecticides

When using Mero with an authorised insecticide, the statutory conditions of use of the plant protection product must be followed. Insect control is not obtained by using Mero alone. Do not use Mero later than the growth stages indicated.

Store well away from seeds, fertilisers and animal feedingstuffs in a safe, dry place designated as an agrochemical store.

PROTECT FROM FROST.

® Mero is a registered Trade Mark of Oleon
© Bayer CropScience Limited 2022

MERO

UFI: E660-M0FM-P00N-FG11

Contains 81.4 % w/w oil (rapeseed fatty acid esters) (EAC1) and ethoxy (7) tridecanol.

Wear protective gloves/protective clothing/eye protection/face protection.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

Plant protection product	Crops			
All authorised herbicides	Forage and grain maize, cereals, sugar beet, ornamental plant production, amenity vegetation, natural surfaces not intended to bear vegetation, permeable surfaces overlying soil, and industrial and amenity areas			
All authorised insecticides	Brussels sprouts, cabbage, broccoli/calabrese, cauliflower and sugar beet			
Maximum concentration (% spray solution):	1.0			
Maximum number of treatments:	Follow the statutory conditions of use of the plant protection product.			
Latest time of application:	Follow the statutory conditions of use of the plant protection product, and: Maize: up to and including GS 53 Cereals: up to and including GS 52 Sugar beet: up to and including 6 leaves unfolded (GS 16) Brussels sprouts: up to and including lateral buds begin to develop (GS 41) Cabbage (Head): Up to heads begin to form (GS 41) Cabbage (leaf): Up to 10% of the leaf mass reached (GS 41) Cauliflower, broccoli/calabrese: Up to and including heads begin to form, growing tip up to 1 cm ² (GS 41) Non-edible crops and non-crop production: follow the statutory conditions of use of the plant protection product			
Other specific restrictions:	Follow all operator protection instructions for the plant protection product. Where the personal protective equipment specified on the adjuvant label is different to that specified on the plant protection product label, the highest level of protective equipment must be worn.			



To access the **Safety Data Sheet** for this product scan the code or use the link below:

www.cropscience.bayer.co.uk/merosds

or alternatively contact your supplier

Bayer CropScience Limited 230, Cambridge Science Park Milton Road, Cambridge CB4 0WB

Telephone: 01223 226500

