

## Emergency Response Preparedness

	<b>Dr. CART</b>	<b>RRT Level 1/2</b>	<b>Anesthesia</b>	<b>Dr. Airway</b>	<b>Stroke</b>	<b>Dr. Stork</b>
<b>Call From Any Hospital Phone</b>	<b>1-4-7</b> (anyone can activate)	<b>1-4-7</b> (anyone can activate)	<b>1-4-7</b> (anyone can activate)	<b>1-4-7</b> (anyone can activate)	<b>1-4-4</b> (RN/MD activates)	<b>1-4-7</b> (anyone can activate)
<b>Overhead Announcement</b>	<b>Yes</b>	<b>No</b>	<b>Yes</b>	<b>Yes</b>	<b>No</b>	<b>Yes</b>
<b>Response Time</b>	<b>Immediate</b>	<b>15 Minutes</b>	<b>Immediate</b>	<b>Immediate</b>	<b>Immediate</b>	<b>Immediate</b>
<b>Clinical Situation</b>	Life-threatening emergency requiring the immediate response of the entire cardio-pulmonary arrest resuscitation team. <i>If unsure, it is preferable to call a Dr. CART</i>	Clinically unstable or deteriorating. Promote better patient outcome by bringing critical care expertise to the bedside	Patient that needs intubation and/or immediate airway evaluation for routine airways	Emergent difficult airway needs and/or emergent ENT needs	New onset of stroke-like symptoms ( $\leq 6$ hours) using the BE-FAST acronym. <i>Ensure: "last known well" and new POC blood glucose</i>	Rapid assessment from Obstetric and Pediatric teams for any obstetric emergency (i.e. precipitous delivery, eclamptic seizure, trauma, cardiac arrest)
<b>Team Members</b>	-Dr. CART (CCD-Cardiology resident on-call/Mitchell-Hospitalist) -Critical Care Nurse -Respiratory Therapy -Anesthesia -Pharmacy -Chaplain -Security -Patient transport -Surgery Resident (page 30134)	-Primary Service (First Contact Provider) -CCON (critical care outreach nurse) -Primary RN -Respiratory Therapy -Patient Transport (DCAM only)	-Anesthesia -Respiratory Therapy -Pharmacy -CCON (non-critical care areas)	-Anesthesia -ENT/Trauma -Surgery -CCON (non-critical care areas) -Respiratory Therapy -Pharmacy	-Primary RN -Neuro Team -First Contact Provider -CCON (non-critical care areas)	-OB Team -NICU Code Blue Team
<b>Coverage</b>	All clinical and non-clinical settings	Non-critical care clinical settings	All clinical and non-clinical settings	All clinical and non-clinical settings	All clinical and non-clinical settings	All clinical and non-clinical settings
<b>Transport Post-Event</b>	All team members transport to ICU after successful resuscitation	CCON, Primary RN, ICU MD transport to ICU	Transport to ICU if needed	Transport to ICU if needed	Transport to CT within 20 minutes	Team members transport to Family Birth Center

## Emergency Response Preparedness

<b>RRT Level 1</b>	<b>RRT Level 2</b>
First Contact Provider	Dr. CART Physician (See PC-44)
Critical Cart Outreach Nurse (CCON)	Critical Care Outreach Nurse (CCON)
Respiratory Therapy	Respiratory Therapy
Transportation Services (DCAM Only)	Transportation Services
Primary Nurse	Security
RRT Pharmacist On-Call (Optional)	RRT Pharmacist On-Call (Optional)

		<b>Routine</b> (Change in condition/Needs medical assistance with response required >15 min)	<b>Urgent</b> (Response required ≤ 15 min) physiologically unstable/Deteriorating)	<b>Emergent</b> (Life-threatening/ Impending or actual loss of pulse/Needs immediate response)
<b>Clinical setting</b>	<b>Patient</b>	Primary Svc	RRT Level 1	Dr. CART
	<b>Visitor/Employee</b>	RRT Level 2	RRT Level 2	Dr. CART
<b>Non-clinical setting</b>	<b>Patient</b>	RRT Level 2	RRT Level 2	Dr. CART
	<b>Visitor/Employee</b>	RRT Level 2	RRT Level 2	Dr. CART

## Emergency Response Preparedness

