

## Drug Facts

### Active ingredients (in each 30 mL dose cup)

Acetaminophen 650 mg. . . . .	Pain reliever/fever reducer
Chlorpheniramine maleate 4 mg. . . . .	Antihistamine
Dextromethorphan HBr 30 mg. . . . .	Cough suppressant

### Purpose

**Uses** temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

### Warnings

**Liver warning:** This product contains acetaminophen.

Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have** • liver disease • glaucoma

- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - cough comes back, or occurs with rash or headache that lasts.
- These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** • take only as directed • use dose cup or tablespoon (Tbsp)

- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL (2 Tbsp) every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

### Other information

- each 30 mL dose cup contains: potassium 7 mg, sodium 45 mg
- store at room temperature

**Inactive ingredients** acesulfame potassium, carboxymethylcellulose sodium, citric acid, FD&C Red No. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium citrate

**Questions?** 1-800-362-1683 [www.vicks.com](http://www.vicks.com)