

## NyQuil™ Cold & Flu HBP

<b>Drug Facts</b>	
<b>Active ingredients (in each 30 mL)</b>	<b>Purpose</b>
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg.....	Cough suppressant
Doxylamine succinate 12.5 mg.....	Antihistamine
<b>Uses</b> temporarily relieves common cold/flu symptoms: <ul style="list-style-type: none"> <li>• cough due to minor throat and bronchial irritation</li> <li>• sore throat</li> <li>• headache</li> <li>• minor aches and pains</li> <li>• fever</li> <li>• runny nose and sneezing</li> </ul>	
<b>Warnings</b> <b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take <ul style="list-style-type: none"> <li>• more than 4 doses in 24 hours, which is the maximum daily amount for this product</li> <li>• with other drugs containing acetaminophen</li> <li>• 3 or more alcoholic drinks every day while using this product</li> </ul> <b>Allergy Alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> <li>• Skin reddening</li> <li>• Blisters</li> <li>• Rash</li> </ul> If a skin reaction occurs, stop use and seek medical help right away <b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
<b>Do not use</b> <ul style="list-style-type: none"> <li>• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>• If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> </ul>	
<b>Ask a doctor before use if you have</b> <ul style="list-style-type: none"> <li>• liver disease</li> <li>• glaucoma</li> <li>• cough that occurs with too much phlegm (mucus)</li> <li>• a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema</li> </ul>	

<b>Drug Facts</b> (continued)	
<ul style="list-style-type: none"> <li>• trouble urinating due to enlarged prostate gland</li> <li>• a sodium-restricted diet</li> </ul>	
<b>Ask a doctor or pharmacist before use if you are</b> <ul style="list-style-type: none"> <li>• taking sedatives or tranquilizers</li> <li>• taking the blood thinning drug warfarin</li> </ul>	
<b>When using this product</b> <ul style="list-style-type: none"> <li>• excitability may occur, especially in children</li> <li>• marked drowsiness may occur</li> <li>• avoid alcoholic drinks</li> <li>• be careful when driving a motor vehicle or operating machinery</li> <li>• alcohol, sedatives, and tranquilizers may increase drowsiness</li> </ul>	
<b>Stop use and ask a doctor if</b> <ul style="list-style-type: none"> <li>• pain or cough gets worse or lasts more than 7 days</li> <li>• fever gets worse or lasts more than 3 days</li> <li>• redness or swelling is present</li> <li>• new symptoms occur</li> <li>• cough comes back or occurs with rash or headache that lasts.</li> </ul> These could be signs of a serious condition.	
<b>If pregnant or breast-feeding</b> , ask a health professional before use. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
<b>Directions</b> <ul style="list-style-type: none"> <li>• take only as directed</li> <li>• only use the dose cup provided</li> <li>• do not exceed 4 doses per 24 hrs</li> </ul>	
adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use
<b>Other information</b> <ul style="list-style-type: none"> <li>• <b>each 30 mL contains:</b> potassium 5 mg, sodium 37 mg</li> <li>• store at room temperature and do not refrigerate.</li> </ul>	
<b>Inactive Ingredients</b> acesulfame potassium, alcohol, citric acid, FD&C Blue No.1, FD&C Red No. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin, sodium, sodium citrate	
<b>Questions? 1-800-362-1683</b>	
<b>Also sold as NyQuil Cold &amp; Flu</b> <a href="http://www.vicks.com">www.vicks.com</a>	