

## Drug Facts

### Active ingredients (in each LiquiCap) Purpose

Acetaminophen 325 mg . . . . . Pain reliever/fever reducer  
 Dextromethorphan HBr 10 mg . . . . . Cough suppressant  
 Phenylephrine HCl 5 mg . . . . . Nasal decongestant

**Uses** temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

### Warnings

**Liver warning:** This product contains acetaminophen.

Severe liver damage may occur if you take

- more than 4 doses in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** • with any other drug containing acetaminophen (prescription or nonprescription).

If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have** • liver disease • heart disease • high blood pressure

- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin.

**When using this product, do not use more than directed.**

**Stop use and ask a doctor if** • you get nervous, dizzy or sleepless

- pain, nasal congestion or cough get worse or last more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

### Directions

- take only as directed
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	2 LiquiCaps with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

### Other information

- store at room temperature

**Inactive ingredients** FD&C Red No. 40, FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide