

DayQuil™ Cold & Flu HBP

Drug Facts	
Active ingredients (in each LiquiCap)	Purpose
Acetaminophen 325 mg.....	Pain reliever/Fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Uses temporarily relieves common cold/flu symptoms: <ul style="list-style-type: none"> • cough due to minor throat & bronchial irritation • sore throat • headache • minor aches & pains • fever 	
Warnings Liver Warning: This product contains acetaminophen. Severe liver damage may occur if you take <ul style="list-style-type: none"> • more than 4 doses in 24 hrs, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> • Skin reddening • Blisters • Rash If a skin reactions occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
Do not use <ul style="list-style-type: none"> • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. <ul style="list-style-type: none"> • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. 	
Ask a doctor before use if you have <ul style="list-style-type: none"> • liver disease • cough that occurs with too much phlegm (mucus) • persistent or chronic cough as occurs with smoking, asthma, or emphysema 	
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.	
Stop use and ask a doctor if <ul style="list-style-type: none"> • pain, nasal congestion or cough get worse or last more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.	

Drug Facts (continued)	
Directions <ul style="list-style-type: none"> • take only as directed • do not exceed 4 doses per 24 hrs 	
adults & children 12 yrs & over	2 LiquiCaps with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Drug Facts (continued)	
Other information <ul style="list-style-type: none"> • store at room temperature 	
Inactive ingredients FD&C Red No. 40, FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide	