

DayQuil Severe Cold and Flu Relief Liquid

PDR P-129116 | POA 010

TAMPER EVIDENT: Do not use if printed shrinkband is missing or broken. Failure to follow these warnings could result in serious consequences.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Drug Facts

Active ingredients (in each 15 mL tablespoon) Purpose
Acetaminophen 325 mg Pain reliever/fever reducer
 Dextromethorphan HBr 10 mg Cough suppressant
 Guaifenesin 200 mg Expectorant
 Phenylephrine HCl 5 mg Nasal decongestant

Uses • temporarily relieves common cold/flu symptoms:
 • nasal congestion • sinus congestion & pressure
 • cough due to minor throat & bronchial irritation • minor aches & pains
 • headache • fever • sore throat • reduces swelling of nasal passages
 • temporarily restores freer breathing through the nose
 • promotes nasal and/or sinus drainage
 • helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hrs
 - child takes more than 4 doses (15 mL each) in 24 hrs
 - taken with other drugs containing acetaminophen
 - adult has 3 or more alcoholic drinks every day while using this product
- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • liver disease • heart disease
 • high blood pressure • thyroid disease • diabetes
 • trouble urinating due to enlarged prostate gland
 • cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if • you get nervous, dizzy or sleepless

- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present • new symptoms occur
 - cough comes back or occurs with rash or headache that lasts.
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see Overdose warning
- use dose cup or tablespoon (TBSP) • do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL (2 TBSP) every 4 hrs
children 6 to under 12 yrs	15 mL (1 TBSP) every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

- when using other DayQuil or NyQuil® products, carefully read each label to ensure correct dosing

Other information

- each 15 mL tablespoon contains: sodium 47 mg
- store at room temperature

Inactive ingredients citric acid, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

Questions? 1-800-251-3374 www.vicks.com



NyQuil Severe Cold and Flu Relief Liquid

PDR P-124021 | POA 021

TAMPER EVIDENT: Do not use if printed shrinkband is missing or broken. Failure to follow these warnings could result in serious consequences.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Drug Facts

Active ingredients (in each 30 mL dose cup) Purpose
Acetaminophen 650 mg Pain reliever/fever reducer
 Dextromethorphan HBr 20 mg Cough suppressant
 Doxylamine succinate 12.5 mg Antihistamine
 Phenylephrine HCl 10 mg Nasal decongestant

Uses • temporarily relieves common cold/flu symptoms:
 • sinus congestion & pressure • cough due to minor throat & bronchial irritation
 • cough to help you sleep • minor aches & pains • headache • fever
 • sore throat • runny nose & sneezing • reduces swelling of nasal passages
 • temporarily restores freer breathing through the nose
 • promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 • to make a child sleep

Ask a doctor before use if you have • liver disease • heart disease
 • high blood pressure • thyroid disease • diabetes • glaucoma
 • cough that occurs with too much phlegm (mucus)
 • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
 • trouble urinating due to enlarged prostate gland • a sodium-restricted diet

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers • taking the blood thinning drug warfarin

When using this product • do not use more than directed

- excitability may occur, especially in children
- marked drowsiness may occur • avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if • you get nervous, dizzy or sleepless

- pain, nasal congestion, or cough gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present • new symptoms occur
 - cough comes back or occurs with rash or headache that lasts.
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see Overdose warning
- use dose cup or tablespoon (TBSP) • do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL (2 TBSP) every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

- when using other DayQuil® or NyQuil products, carefully read each label to ensure correct dosing

Other information

- each 30 mL dose cup contains: sodium 81 mg
- store at room temperature

Inactive ingredients alcohol, citric acid, D&C Yellow No. 10, FD&C Yellow No. 6, FD&C Green No. 3, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose

Questions? 1-800-362-1683 www.vicks.com

