

<b>Drug Facts</b> NyQuil Cold & Flu	
<b>Active ingredients (in each 30 mL dose cup)</b>	<b>Purpose</b>
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine
<b>Uses</b> temporarily relieves common cold/flu symptoms: <ul style="list-style-type: none"> <li>cough due to minor throat &amp; bronchial irritation</li> <li>sore throat</li> <li>headache</li> <li>minor aches &amp; pains</li> <li>fever</li> <li>runny nose &amp; sneezing</li> </ul>	
<b>Warnings</b> <b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take <ul style="list-style-type: none"> <li>more than 4 doses in 24 hrs, which is the maximum daily amount for this product</li> <li>with other drugs containing acetaminophen</li> <li>3 or more alcoholic drinks every day while using this product</li> </ul> <b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
<b>Do not use</b> <ul style="list-style-type: none"> <li>with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> </ul>	
<b>Ask a doctor before use if you have</b> <ul style="list-style-type: none"> <li>liver disease</li> <li>glaucoma</li> <li>cough that occurs with too much phlegm (mucus)</li> <li>a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema</li> <li>trouble urinating due to enlarged prostate gland</li> <li>a sodium-restricted diet</li> </ul>	
<b>Ask a doctor or pharmacist before use if you are</b> <ul style="list-style-type: none"> <li>taking sedatives or tranquilizers</li> <li>taking the blood thinning drug warfarin</li> </ul>	
<b>When using this product</b> <ul style="list-style-type: none"> <li>excitability may occur, especially in children</li> <li>marked drowsiness may occur</li> <li>avoid alcoholic drinks</li> <li>be careful when driving a motor vehicle or operating machinery</li> <li>alcohol, sedatives, &amp; tranquilizers may increase drowsiness</li> </ul>	
<b>Stop use and ask a doctor if</b> <ul style="list-style-type: none"> <li>pain or cough gets worse or lasts more than 7 days</li> <li>fever gets worse or lasts more than 3 days</li> <li>redness or swelling is present</li> <li>new symptoms occur</li> <li>cough comes back or occurs with rash or headache that lasts.</li> </ul> These could be signs of a serious condition.	
<b>If pregnant or breast-feeding</b> , ask a health professional before use.	

<b>Drug Facts</b> DayQuil Cold & Flu	
<b>Active ingredients (in each 15 mL tablespoon)</b>	<b>Purpose</b>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant
<b>Uses</b> temporarily relieves common cold/flu symptoms: <ul style="list-style-type: none"> <li>nasal congestion</li> <li>cough due to minor throat &amp; bronchial irritation</li> <li>sore throat</li> <li>headache</li> <li>minor aches &amp; pains</li> <li>fever</li> </ul>	
<b>Warnings</b> <b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if <ul style="list-style-type: none"> <li>adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product</li> <li>child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product</li> <li>taken with other drugs containing acetaminophen</li> <li>adult has 3 or more alcoholic drinks every day while using this product</li> </ul> <b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
<b>Do not use</b> <ul style="list-style-type: none"> <li>with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> </ul>	
<b>Ask a doctor before use if you have</b> <ul style="list-style-type: none"> <li>liver disease</li> <li>heart disease</li> <li>high blood pressure</li> <li>thyroid disease</li> <li>diabetes</li> <li>trouble urinating due to enlarged prostate gland</li> <li>cough that occurs with too much phlegm (mucus)</li> <li>persistent or chronic cough such as occurs with smoking, asthma, or emphysema</li> <li>a sodium restricted diet</li> </ul>	
<b>Ask a doctor or pharmacist before use if you are</b> taking the blood thinning drug warfarin.	
<b>When using this product, do not use more than directed.</b>	
<b>Stop use and ask a doctor if</b> <ul style="list-style-type: none"> <li>you get nervous, dizzy or sleepless</li> <li>pain, nasal congestion or cough get worse or last more than 5 days (children) or 7 days (adults)</li> <li>fever gets worse or lasts more than 3 days</li> <li>redness or swelling is present</li> <li>new symptoms occur</li> <li>cough comes back, or occurs with rash or headache that lasts.</li> </ul> These could be signs of a serious condition.	
<b>If pregnant or breast-feeding</b> , ask a health professional before use.	

<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.							
<b>Directions</b> <ul style="list-style-type: none"> <li>take only as directed</li> <li>use dose cup or tablespoon (TBSP)</li> <li>do not exceed 4 doses per 24 hours</li> </ul> <table border="1"> <tr> <td>adults &amp; children 12 yrs &amp; over</td> <td>30 mL (2 TBSP) every 6 hrs</td> </tr> <tr> <td>children 4 to under 12 yrs</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 yrs</td> <td>do not use</td> </tr> </table>		adults & children 12 yrs & over	30 mL (2 TBSP) every 6 hrs	children 4 to under 12 yrs	ask a doctor	children under 4 yrs	do not use
adults & children 12 yrs & over	30 mL (2 TBSP) every 6 hrs						
children 4 to under 12 yrs	ask a doctor						
children under 4 yrs	do not use						
<b>Other information</b> <ul style="list-style-type: none"> <li>each 30 mL dose cup contains: potassium 5 mg, sodium 37 mg</li> <li>store at room temperature</li> </ul>							
<b>Inactive ingredients</b> acesulfame potassium, alcohol, citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Yellow No. 6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate							
<b>Questions?</b> 1-800-362-1683							

<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.									
<b>Directions</b> <ul style="list-style-type: none"> <li>take only as directed</li> <li>use dose cup or tablespoon (TBSP)</li> <li>do not exceed 4 doses per 24 hrs</li> </ul> <table border="1"> <tr> <td>adults &amp; children 12 yrs &amp; over</td> <td>30 mL (2 TBSP) every 4 hrs</td> </tr> <tr> <td>children 6 to under 12 yrs</td> <td>15 mL (1 TBSP) every 4 hrs</td> </tr> <tr> <td>children 4 to under 6 yrs</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 yrs</td> <td>do not use</td> </tr> </table>		adults & children 12 yrs & over	30 mL (2 TBSP) every 4 hrs	children 6 to under 12 yrs	15 mL (1 TBSP) every 4 hrs	children 4 to under 6 yrs	ask a doctor	children under 4 yrs	do not use
adults & children 12 yrs & over	30 mL (2 TBSP) every 4 hrs								
children 6 to under 12 yrs	15 mL (1 TBSP) every 4 hrs								
children 4 to under 6 yrs	ask a doctor								
children under 4 yrs	do not use								
<b>Other information</b> <ul style="list-style-type: none"> <li>each 15 mL tablespoon contains: sodium 46 mg</li> <li>store at room temperature</li> </ul>									
<b>Inactive ingredients</b> citric acid, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum									
<b>Questions?</b> 1-800-251-3374									