

## This material is designed to be used by patients who have been prescribed sublingual apomorphine

KYNMOBI® is a medicine to place under the tongue (sublingual film) that contains the active substance apomorphine hydrochloride. It is for use, as needed, with other Parkinson's medicines taken orally (by your mouth) to reduce the amount of time spent in an "OFF", a period during the day when your Parkinson's symptoms are noticeably worse. Parkinson's disease is a progressive disease of the nervous system that causes shaking and affects your movement

### HOW LONG DOES IT TAKE TO ACT?



Improvements should be noticed after 15 to 30 minutes, but may vary from patient to patient.

### HOW TO STORE SUBLINGUAL APOMORPHINE AT HOME?



Keep the apomorphine film in the sachet until ready to take it.



Do not store at temperatures above 25°C. Store in the sachet to protect from light and moisture.



Are you noticing you need this more? Contact your clinician or healthcare team.

### HOW CAN I IMPROVE MY QUALITY OF LIFE?

There are 3 fundamental factors to keep yourself active everyday:



Nourishment



Exercise



Health and Mental Activity



## Check out our **bialive** to learn more about Parkinson's Disease

This resource has been developed by Bial.

# Parkinson's Disease and KYNMOBI®

## THE IMPORTANCE OF TAKING CARE OF YOUR MOUTH

This material is intended for patients who have been prescribed KYNMOBI® (apomorphine hydrochloride)



**The medicine you are about to take, may cause some discomfort. Most of these effects are mild to moderate and can usually be reduced with general care.**

**Before placing the medicine in your mouth, you should:**

- Moisten your mouth (for example, by rinsing with water or a saline solution)

**When using the medicine, do not swallow until it has completely dissolved.**

**After using the medicine, you should:**

- Rinse your mouth again after the medicine has been dispersed and absorbed
- This is about 3-5 minutes after it has been absorbed



- Remove them before using the medicine
- Remove any excess denture adhesive
- Avoid applying denture adhesive straight after using the medicine

**Mouth irritation is a very common side effect of Kynmobi. You should tell your doctor if you develop any of the following signs or symptoms:¹**

- Redness
- Mouth sores (ulceration)
- Dryness of the mouth, lips or tongue
- Swelling
- Mouth pain

These signs and symptoms may go away if Kynmobi is stopped. Speak with your clinician if you experience any of these changes.



**Maintain good oral hygiene and moisten your mouth before taking KYNMOBI®**

If you experience side effects talk to your doctor or other healthcare professional. This includes any possible side effects not listed in the package leaflet. To report a side effect or product complaint with a Bial product please call Bial on +44 (0) 1753 916 010 or [GCS-Safetybial.UKIE@cencora.com](mailto:GCS-Safetybial.UKIE@cencora.com).

Additionally, reporting forms and further information can be found at <https://yellowcard.mhra.gov.uk/> or search for MHRA Yellow Card in the Google Play or Apple App Store or in Ireland via HPRA pharmacovigilance website [www.hpra.ie](http://www.hpra.ie)

By reporting side effects you can help provide more information on the safety of medicines.

1. KYNMOBI® (Apomorphine hydrochloride). Summary of Product Characteristics. Bial Pharma UK Ltd. Available at: <https://www.medicines.org.uk/>. Access date - August 2026

**Bial** Keeping life in mind

**KYNMOBI®**  
(apomorphine HCl) sublingual film  
10 mg • 15 mg • 20 mg • 25 mg • 30 mg