

ConnectAD™



Welcome to **ConnectAD**TM, a clinical case series created by the Eli Lilly and Company Neuroscience medical education team. This series is intended to connect healthcare professionals to resources that help them detect, diagnose, and manage Alzheimer's disease.




Disclaimer

The content for this clinical case was developed in collaboration between a group of global clinicians who care for patients with Alzheimer's disease and Eli Lilly and Company.

A variety of cognitive and diagnostic tests can reasonably be used in the detection and diagnosis of Alzheimer's disease. Inclusion of specific cognitive and/or diagnostic tests in this case reflects the diversity of clinical preferences, and the use of particular diagnostic tools does not imply endorsement or recommendation by Lilly.

Learning Objectives

- Through completing this course, you will have a deeper understanding of:
 - The clinical presentation of Alzheimer's disease
 - How to integrate clinical and biomarker assessments to make an accurate diagnosis of Alzheimer's disease in the earliest stages



Connect with Maria



Our Patient

Maria



72 years old



Female



Hispanic



Married, 4 children, 3 grandchildren



College degree



Accountant (retired)



Family history

- Cardiovascular disease
- Type 2 diabetes
- Alzheimer's disease (mother)



Clinical Information and History

Maria has experienced onset of the following symptoms, which have gradually progressed over the last 2 years:

- Difficulty remembering dates and events
- Problems finding words



Clinical history

- Type 2 diabetes
- Hypertension




General health

- Heart rate: 72 bpm
- Blood pressure: 138/70 mmHg



Current medication (class)

- Angiotensin-converting enzyme (ACE) inhibitor
- Biguanide



Initial Clinical Assessment



General neurological exam: Normal



Mental status

- Alert, insightful, good effort on testing
- Activities of daily living: Independent



Cognition

- MoCA: 25/30 (normal ≥ 26)
 - Word recall: 1/5
 - Figure copy: Incorrect
 - Clock drawing: 3/3
- BNT Short Form: 12/15 with circumlocutions (normal ≥ 12)



Additional Findings



Blood work within normal limits, including:

- Complete blood count (CBC)
- Electrolytes
- Glucose
- Creatinine
- Thyroid stimulating hormone (TSH)
- Vitamin B12



Brain MRI

- Mild diffuse cortical atrophy
- Mild microangiopathy
- Small old lacune in left external capsule



CSF assay

- $A\beta_{42}/A\beta_{40}$: Low
- P-tau: High
- T-tau: High



Given the patient information presented, what is the diagnosis?

1. Generalized anxiety
2. MCI due to AD
3. Preclinical AD
4. Vascular cognitive impairment
5. Other



***Given the patient information presented,
what is the diagnosis?***

Option 2:
MCI due to AD



Justification for Diagnosis

Why is Maria diagnosed with MCI due to AD?

Maria is presenting with clinical symptoms consistent with the earliest stages of AD, which are corroborated by clinical assessment.

Her MRI results suggest a mild vascular contribution to her symptoms, but are not definitive, which can be typical for someone of her age and early clinical presentation.

However, the results of her CSF analysis confirm the presence of AD pathology. Specifically, the decreased concentration of $A\beta_{42}$ indicates amyloidosis, while the elevation in tau species indicates axonal damage, and suggests the presence of neurofibrillary tangles.¹



Key Learnings in Maria's Case (1 of 3)

MCI due to AD is vastly underdiagnosed,¹ and can be difficult to distinguish from other causes of cognitive impairment.² AD biomarkers can provide the confirmatory information needed for diagnosis³

- Despite variations in labeling, there is broad agreement on the stages and continuum of events in the progression of AD⁴
- Evidence of AD pathology is present many years before cognitive symptoms appear. This period is referred to as the 'preclinical' period⁴
- As the disease progresses, patients will experience increasing cognitive impairment, which may impact ability to perform activities of daily living. These sequential stages of disease are referred to as 'mild cognitive impairment', or 'dementia (characterized as 'mild', 'moderate', or 'severe') due to AD'⁴



AD=Alzheimer's Disease.

1. Liu Y, et al. *J Prev Alzheimers Dis.* 2024;11(1):7-12. 2. Albert MS, et al. *Alzheimers Dement.* 2011;7(3):270-279.

3. <https://alz.org/media/Documents/scientific-conferences/Figures-and-Tables-Clinical-Criteria-for-Staging-and-Diagnosis-for-Public-Comment-Draft-2.pdf>. (Accessed January 2024).

4. Porsteinsson AP, et al. *J Prev Alz Dis.* 2021;3(8):371-386.

Key Learnings in Maria's Case (2 of 3)

The CSF biomarker assay can be used to confirm the presence of pathology consistent with AD¹

CSF assays

- Quantitatively measure the levels of A β and tau protein within the fluid of the lumbar sac²
- Reflect the rates of A β and tau protein production and clearance²
- Advantages include:
 - Less expensive than PET by 10-15-fold³
 - Simultaneous information on A β and tau biomarkers⁴
- Limitations include:
 - CSF is obtained via lumbar puncture; this is invasive and can be uncomfortable for patients⁵
 - Limited availability outside of specialized clinics⁶
 - Do not detect regional A β or tau deposition^{2,3}



A β =Amyloid Beta; AD=Alzheimer's Disease; CSF=Cerebrospinal Fluid; MCI=Mild Cognitive Impairment; PET=Positron Emission Tomography.

1. <https://alz.org/media/Documents/scientific-conferences/Figures-and-Tables-Clinical-Criteria-for-Staging-and-Diagnosis-for-Public-Comment-Draft-2.pdf>. (Accessed January 2024). 2. Jack CR Jr, et al. *Alzheimers Dement*. 2018;14:535-562.

3. Hansson O, et al. *Alzheimer's Res Ther*. 2019;11(1):34. 4. Dubois B, et al. *Lancet Neurol*. 2021;20(6):484-496. 5. Lee JC, et al. *Exp Mol Med*. 2019;51(5):1-10. 6. Zetterberg H, et al. *Alzheimer's Dement (Amst)*. 2019;784-786.

Key Learnings in Maria's Case (3 of 3)

MRI can be a useful assessment to¹:

- Rule out non-AD conditions that can cause cognitive decline
- Provide information suggestive of AD; for example, observed hippocampal atrophy

However, the scan does not provide information on amyloid and tau biomarkers, and cannot be used to diagnose AD in isolation²

Structural neuroimaging (MRI)

- Can detect atrophy, which correlates with cognitive status and is a marker of neurodegeneration¹
- Advantages include:
 - Noninvasive and widely available³
 - No ionizing irradiation^{4,5}
 - Excellent soft tissue contrast and high spatial resolution^{4,5}
- Disadvantages include:
 - Some patients find the scanner claustrophobic^{5,6}
 - Patients with magnetic metal implants should not receive MRI exams⁵
 - Atrophy patterns seen are not specific to AD⁶




AD=Alzheimer's Disease; MRI=Magnetic Resonance Imaging.

1. Park M, Moon WJ. *Korean J Radiol.* 2016;17(6):827-845. 2. Porsteinsson AP, et al. *J Prev Alz Dis.* 2021;3(8):371-386. 3. McEvoy LK, Brewer JB. *Expert Rev Neurother.* 2010;10(11):1675-1688.

4. Pysz MA, et al. *Clin Radiol.* 2010;65(7):500-516. 5. <https://www.fda.gov/radiation-emitting-products/mri-magnetic-resonance-imaging/benefits-and-risks>. (Accessed January 2024).

6. Johnson KA, et al. *Cold Spring Harb Perspect Med.* 2012;2:a006213.



Thank you for connecting
with Maria

